arXiv:math-ph/0604028v1 13 Apr 2006

Analysis on q-deformed quantum spaces

Hartmut Wachter*

Max-Planck-Institute
for Mathematics in the Sciences
Inselstr. 22, D-04103 Leipzig

Arnold-Sommerfeld-Center
Ludwig-Maximilians-Universität
Theresienstr. 37, D-80333 München

Abstract

A q-deformed version of classical analysis is given to quantum spaces of physical importance, i.e. Manin plane, q-deformed Euclidean space in three or four dimensions, and q-deformed Minkowski space. The subject is presented in a rather complete and selfcontained way. All relevant notions are introduced and explained in detail. The different possibilities to realize the objects of q-deformed analysis are discussed and their elementary properties are studied. In this manner attention is focused on star products, q-deformed tensor products, q-deformed translations, q-deformed partial derivatives, dual pairings, q-deformed exponentials, and q-deformed integration. The main concern of this work is to show that these objects fit together in a consistent framework, which is suitable to formulate physical theories on quantum spaces.

*e-mail: Hartmut.Wachter@physik.uni-muenchen.de

Contents

1 Introduction **3**
1.1 General motivation for deforming spacetime 3
1.2 q-Deformation of symmetries as an attempt to get a more detailed description of nature
1.3 Foundations of q-deformed analysis 4
1.4 Intention and content of this work 5

2 Star products, q-tensor products, and q-translations **6**
2.1 Definition of star products and q-deformed tensor products . . 6
2.2 Definition and basic properties of q-deformed translations . . . 12
2.3 Conjugation properties and crossing symmetries 19

3 Partial derivatives on quantum spaces **26**
3.1 Leibniz rules and representations of partial derivatives 26
3.2 Conjugation properties and crossing symmetries 30
3.3 Further correspondences between actions of partial derivatives 33

4 Dual pairings and q-exponentials **35**
4.1 Definition and basic properties of dual pairings 36
4.2 Definition and basic properties of q-deformed exponentials . . 39
4.3 q-Deformed analogs of Taylor rules 49
4.4 Conjugation properties and crossing symmetries 51

5 Integration on quantum spaces **54**
5.1 Inverse partial derivatives and their representations 54
5.2 Basic properties of q-deformed integrals 60
5.3 Integrals over the whole space and their translation invariance 67
5.4 Conjugation properties of q-deformed integrals 79

6 Conclusion **81**

A Quantum spaces **82**

B Integrals for q-deformed Minkowski space **84**

1 Introduction

1.1 General motivation for deforming spacetime

Relativistic quantum field theory is not a fundamental theory, since its formalism leads to divergencies. In some cases like that of quantum electrodynamics one is able to overcome the difficulties with the divergencies by applying the so-called renormalization procedure due to Feynman, Schwinger, and Tomonaga [1]. Unfortunately, this procedure is not successful if we want to deal with quantum gravity. Despite the fact that gravitation is a rather weak interaction we are not able to treat it perturbatively. The reason for this lies in the fact that transition amplitudes of n-th order to the gravitation constant diverge like a momentum integral of the general form [2]

$$\int p^{2n-1} dp, \tag{1}$$

leaving us with an infinite number of ultraviolet divergent Feynman diagrams that cannot be removed by redefining finitely many physical parameters.

It is surely legitimate to ask for the reason for these fundamental difficulties. It is commonplace that the problems with the divergences in relativistic quantum field theory result from an incomplete description of spacetime at very small distances [1]. Niels Bohr and Werner Heisenberg have been the first who suggested that quantum field theories should be formulated on a spacetime lattice [3,4]. Such a spacetime lattice would imply the existence of a smallest distance a with the consequence that plane-waves of wave-length smaller than twice the lattice spacing could not propagate. In accordance with the relationship between wave-length λ and momentum p of a plane-wave, i.e.

$$\lambda \geq \lambda_{\min} = 2a \quad \Rightarrow \quad \frac{1}{\lambda} \sim p \leq p_{\max} \sim \frac{1}{2a}, \tag{2}$$

it follows then that physical momentum space would be bounded. Hence, the domain of all momentum integrals in Eq. (1) would be bounded as well with the consequence that momentum integrals should take on finite values.

1.2 q-Deformation of symmetries as an attempt to get a more detailed description of nature

Discrete spacetime structures in general do not respect classical Poincaré symmetry. A possible way out of this difficulty is to modify not only spacetime but also its corresponding symmetries. How are we to accomplish this? First of all let us recall that classical spacetime symmetries are usually described by Lie groups. If we realize that Lie groups are manifolds the Gelfand-Naimark theorem tells us that Lie groups can be naturally embedded in the category of algebras [5]. The utility of this interrelation lies in formulating the geometrical structure of Lie groups in terms of a Hopf structure [6]. The point is that during the last two decades generic methods have been discovered for continuously deforming matrix groups and Lie algebras within the category of Hopf algebras. It is this development which finally led to the arrival of quantum groups and quantum spaces [7–13].

From a physical point of view the most realistic and interesting deformations are given by q-deformed versions of Minkowski space and Euclidean spaces as well as their corresponding symmetries, i.e. respectively Lorentz symmetry and rotational symmetry [14–18]. Further studies even allowed to establish differential calculi on these q-deformed quantum spaces [19–22] representing nothing other than q-analogs of translational symmetry. In this sense we can say that q-deformations of the complete Euclidean and Poincaré symmetries are now available [23]. Finally, Julius Wess and his coworkers were able to show that q-deformation of spaces and symmetries can indeed lead to the wanted discretizations of the spectra of spacetime observables [24–27], which nourishes the hope that q-deformation might give a new method to regularize quantum field theories [28–31].

1.3 Foundations of q-deformed analysis

In order to formulate quantum field theories on quantum spaces it is necessary to provide us with some essential tools of a q-deformed analysis [32]. The main question is how to define these new tools, which should be q-analogs of classical notions. Towards this end the considerations of Shahn Majid have proved very useful [33–36]. The key idea of this approach is that all the quantum spaces to a given quantum symmetry form a braided tensor category. Consequently, operations and objects concerning quantum spaces must rely on this framework of a braided tensor category, in order to guarantee their

well-defined behavior under quantum group transformations. This so-called principle of covariance can be seen as the essential guideline for constructing a consistent theory.

In our previous work [37–44] we have applied these general considerations, exposed in Refs. [35,36,45–47], to quantum spaces of physical importance, i.e. q-deformed quantum plane, q-deformed Euclidean space with three or four dimensions, and q-deformed Minkowski space. In this manner, we obtained explicit expressions for computing star products, operator representations of partial derivatives and symmetry generators, q-integrals, q-exponentials, q-translations, and braided products. With this toolbox of essential elements of q-analysis we should be able to develop a q-deformed version of quantum field theory along the same line of reasonings as in the classical case. To achieve this we have to gain further insight into the properties of the new objects. In other words, we need a set of consistent rules telling us how to perform calculations with the new objects.

In this respect we have to be aware of one remarkable difference between q-deformed objects and their classical counterparts. For each of our q-deformed objects we can find different realizations, which in the undeformed case become identical. The reason for this lies in the fact that the braided tensor category in which a deformed object lives is not uniquely determined. This becomes quite clear, if one realizes that each braided category is characterized by a so-called braiding mapping Ψ. Its inverse Ψ^{-1} gives an equally good braiding which, in general, leads to a different braided category. But we have to deal with both categories, since they are linked via the operation of conjugation [48–50].

1.4 Intention and content of this work

Important developments in theoretical physics are often accompanied by the arrival of new mathematical techniques. Classical mechanics, for example, is deeply connected to classical analysis. It is also well-known that general relativity and quantum mechanics are heavily based on ideas of Riemannian geometry and functional analysis, respectively. As it was pointed out above, the mathematics of quantum groups and quantum spaces can provide a framework for a more detailed description of spacetime at very small distances. If this is the case the formalism should be prepared in a manner that allows to formulate and evaluate physical theories on quantum spaces in a rather systematical way.

In this article we develop basic ideas of q-analysis on quantum spaces of physical importance in a rather complete and self-contained way. We introduce and explain all relevant notions in detail and discuss the different possibilities for realizing the objects of q-analysis. Furthermore, we give a review of elementary properties and derive useful calculational rules. We show how the various objects fit together in a consistent framework. It is also our aim to introduce a convenient notation, which will prove useful in further investigations. In doing so, we hope to provide the foundations for performing concrete calculations, as they are necessary in formulating and evaluating field theories on quantum spaces.

In particular, we intend to proceed as follows. In Section 2 we deal with star products, q-deformed tensor products, and q-deformed translations. In Section 3 we concern ourselves with q-deformed partial derivatives. Section 4 is devoted to the discussion of dual pairings and q-deformed exponentials. In Section 5 we introduce a powerful concept of integration on q-deformed quantum spaces. Section 6 closes our considerations by a short conclusion. For reference and for the purpose of introducing consistent and convenient notation, we provide a review of key notation and results in Appendix A. Last but not least Appendix B contains a discussion of the details about integration on q-deformed Minkowski space.

2 Star products, q-tensor products, and q-translations

This section is devoted to star products, q-deformed tensor products, and q-deformed translations. Our considerations apply to quantum spaces like the Manin plane, the q-deformed Euclidean space in three or four dimensions, and the q-deformed Minkowski space.

2.1 Definition of star products and q-deformed tensor products

For our purposes, it is at first sufficient to consider a quantum space as an algebra of formal power series in non-commuting coordinates $X^1, X^2, \ldots, X^n$,

i.e.

$$\mathcal{A}_q = \mathbb{C}\left[[X_1, \ldots, X_n]\right]/\mathcal{I}, \tag{3}$$

where $\mathcal{I}$ denotes the ideal generated by the relations of the non-commuting coordinates. The term 'coordinate', however, is a little bit misleading, since the generators of quantum spaces cannot only play the role of coordinates but also that of partial derivatives or differentials.

The two-dimensional Manin plane is one of the simplest examples for a quantum space [51]. It consists of all the power series in two coordinates X^1 and X^2 being subject to the commutation relations

$$X^1 X^2 = q X^2 X^1, \quad q > 1. \tag{4}$$

We can think of q as a deformation parameter measuring the coupling among different spatial degrees of freedom. In the classical case, i.e. if q becomes 1 we regain commutative coordinates.

Next, we would like to focus our attention on the question how to perform calculations on an algebra of quantum space coordinates. This can be accomplished by a kind of pullback transforming operations on non-commutative coordinate algebras to those on commutative ones. For this to become more clear, we have to realize that the non-commutative algebras we are dealing with satisfy the *Poincaré-Birkhoff-Witt property*. It tells us that the dimension of a subspace of homogenous polynomials should be the same as for commuting coordinates. This property is the deeper reason why monomials of a given normal ordering constitute a basis of $\mathcal{A}_q$. Due to this fact, we can establish a vector space isomorphism between $\mathcal{A}_q$ and a commutative algebra $\mathcal{A}$ generated by ordinary coordinates $x^1, x^2, \ldots, x^n$:

$$\begin{gathered} \mathcal{W} : \mathcal{A} \longrightarrow \mathcal{A}_q, \\ \mathcal{W}((x^1)^{i_1} \ldots (x^n)^{i_n}) \equiv (X^1)^{i_1} \ldots (X^n)^{i_n}. \end{gathered} \tag{5}$$

This vector space isomorphism can even be extended to an algebra isomorphism by introducing a non-commutative product in $\mathcal{A}$, the so-called *star product* [52–54]. This product is defined via the relation

$$\mathcal{W}(f \circledast g) = \mathcal{W}(f) \cdot \mathcal{W}(g), \tag{6}$$

being tantamount to

$$f \circledast g \equiv \mathcal{W}^{-1}\left(\mathcal{W}\left(f\right) \cdot \mathcal{W}\left(g\right)\right), \tag{7}$$

where f and g are formal power series in $\mathcal{A}$. In the case of the Manin plane, the star product is of the well-known form

$$\begin{aligned} f(x^i) \circledast g(x^j) &= q^{-\hat{n}_{x^2}\hat{n}_{y^1}} \left[f(x^i)\, g(y^j)\right]\big|_{y\to x} \\ &= f(x^i)\, g(x^j) + O(h), \quad \text{with} \quad q = e^h, \end{aligned} \tag{8}$$

where we have introduced the operators

$$\hat{n}_{x^i} \equiv x^i \frac{\partial}{\partial x^i}, \quad i = 1, 2. \tag{9}$$

From the last equality in (8) we see that star products on quantum spaces lead to modifications of commutative products. Evidently, this modifications vanish in the classical limit $q \to 1$.

Next, we would like to deal with tensor products of quantum spaces. To this end, we have to specify the commutation relations between generators of distinct quantum spaces. These relations are completely determined by the requirement of being consistent with the action of a Hopf algebra $\mathcal{H}$ describing the symmetry of the quantum spaces. Making for the commutation relations between two quantum space generators $X^i \in \mathcal{A}_q$ and $Y^j \in \mathcal{A}'_q$ an ansatz of the form

$$X^i Y^j = k\, C^{ij}_{kl}\, Y^k X^l, \quad k, C^{ij}_{kl} \in \mathbb{C}, \tag{10}$$

the coefficients C^{ij}_{kl} have to be determined in such a way that the following condition of covariance is satisfied:

$$(h_{(1)} \triangleright X^i)(h_{(2)} \triangleright Y^j) = k\, C^{ij}_{kl}\, (h_{(1)} \triangleright Y^k)(h_{(2)} \triangleright X^l), \tag{11}$$

where h denotes an arbitrary element of $\mathcal{H}$. Notice that the coproduct of h is written in the so-called Sweedler notation, i.e. $\Delta(h) = h_{(1)} \otimes h_{(2)}$, and repeated indices are to be summed.

It is well-known that for a quasitriangular Hopf algebra $\mathcal{H}$ the last identity is fulfilled if we substitute for the coefficients C^{ij}_{kl} a suitable representation of the universal R-matrix $\mathcal{R} = \mathcal{R}_{[1]} \otimes \mathcal{R}_{[2]} \in \mathcal{H} \otimes \mathcal{H}$. To be more specific, we

have

$$\begin{aligned} X^iY^j &= (\mathcal{R}_{[2]} \triangleright Y^j)\,(\mathcal{R}_{[1]} \triangleright X^i) \\ &= (Y^j \triangleleft \mathcal{R}_{[2]})\,(X^i \triangleleft \mathcal{R}_{[1]}) = k\,\hat{R}^{ij}_{kl}\,Y^kX^l. \end{aligned} \tag{12}$$

Alternatively, we can take a representation of the transposed inverse of $\mathcal{R}$, i.e. $\tau(\mathcal{R}^{-1}) = \mathcal{R}^{-1}_{[2]} \otimes \mathcal{R}^{-1}_{[1]}$, giving us

$$\begin{aligned} X^iY^j &= (\mathcal{R}^{-1}_{[1]} \triangleright Y^j)\,(\mathcal{R}^{-1}_{[2]} \triangleright X^i) \\ &= (Y^j \triangleleft \mathcal{R}^{-1}_{[1]})\,(X^i \triangleleft \mathcal{R}^{-1}_{[2]}) = k^{-1}(\hat{R}^{-1})^{ij}_{kl}\,Y^kX^l. \end{aligned} \tag{13}$$

To understand the second identity in (12) and (13), one has to realize that

$$\begin{aligned} (S^{-1} \otimes S^{-1}) \circ \mathcal{R} &= \mathcal{R}, \\ (S^{-1} \otimes S^{-1}) \circ \mathcal{R}^{-1} &= \mathcal{R}^{-1}, \end{aligned} \tag{14}$$

and

$$S^{-1}(h) \triangleright a = a \triangleleft h, \quad h \in \mathcal{H}, \quad a \in \mathcal{A}_q, \tag{15}$$

where S^{-1} denotes the inverse of the antipode of $\mathcal{H}$.

The identities in (12) and (13) extend to arbitrary quantum space elements. Especially, we have for the commutation relations of a single quantum space generator X^i with an arbitrary element g of another quantum space

$$\begin{aligned} X^ia &= (\mathcal{R}_{[2]} \triangleright a)\,(\mathcal{R}_{[1]} \triangleright X^i) = \big((\bar{\mathcal{L}}_x)^i_j \triangleright a\big)\,X^j, \\ aX^i &= (X^i \triangleleft \mathcal{R}_{[2]})\,(a \triangleleft \mathcal{R}_{[1]}) = X^j\big(a \triangleleft (\mathcal{L}_x)^i_j\big), \end{aligned} \tag{16}$$

or

$$\begin{aligned} X^ia &= (\mathcal{R}^{-1}_{[1]} \triangleright a)\,(\mathcal{R}^{-1}_{[2]} \triangleright X^i) = \big((\mathcal{L}_x)^i_j \triangleright a\big)\,X^j, \\ aX^i &= (X^i \triangleleft \mathcal{R}^{-1}_{[1]})\,(a \triangleleft \mathcal{R}^{-1}_{[2]}) = X^j\big(a \triangleleft (\bar{\mathcal{L}}_x)^i_j\big), \end{aligned} \tag{17}$$

where we introduced the so-called L-matrix together with its conjugate. It should be obvious that the entries of the L-matrices live in the quantum algebra $\mathcal{H}$. As an example, we give the non-vanishing entries of the L-matrices in the case of the Manin plane [43]:

$$(\mathcal{L}_a)^1_1 = \Lambda(a)\tau^{-1/4},$$

$$
\begin{aligned}
(\mathcal{L}_a)^2_1 &= -q\lambda\Lambda(a)\tau^{-1/4}T^+, \\
(\mathcal{L}_a)^2_2 &= \Lambda(a)\tau^{1/4},
\end{aligned} \tag{18}
$$

and likewise

$$
\begin{aligned}
(\bar{\mathcal{L}}_a)^1_1 &= \Lambda^{-1}(a)\tau^{1/4}, \\
(\bar{\mathcal{L}}_a)^1_2 &= q^{-1}\lambda\Lambda^{-1}(a)\tau^{-1/4}T^-, \\
(\bar{\mathcal{L}}_a)^2_2 &= \Lambda^{-1}(a)\tau^{-1/4}.
\end{aligned} \tag{19}
$$

Notice that $\tau^{\pm 1/4}$ and $T^{\pm}$ denote generators of the quantum algebra $U_q(su_2)$ and $\Lambda(a)$ stands for a unitary scaling operator. The constant λ is given by $\lambda \equiv q - q^{-1}$. The L-matrices for q-deformed Euclidean space in three or four dimensions and those for q-deformed Minkowski space can easily be read off from the results in Refs. [38] and [43].

By virtue of the algebra isomorphism $\mathcal{W}$ we are able to introduce the following operations:

$$
\begin{aligned}
f(x^i) \odot_L g(y^j) &\equiv \mathcal{W}^{-1}\big(\mathcal{R}^{-1}_{[1]} \triangleright \mathcal{W}(g)\big) \otimes \mathcal{W}^{-1}\big(\mathcal{R}^{-1}_{[2]} \triangleright \mathcal{W}(f)\big), \\
f(x^i) \odot_{\bar{R}} g(y^j) &\equiv \mathcal{W}^{-1}\big(\mathcal{W}(g) \triangleleft \mathcal{R}^{-1}_{[1]}\big) \otimes \mathcal{W}^{-1}\big(\mathcal{W}(f) \triangleleft \mathcal{R}^{-1}_{[2]}\big),
\end{aligned} \tag{20}
$$

and

$$
\begin{aligned}
f(x^i) \odot_{\bar{L}} g(y^j) &\equiv \mathcal{W}^{-1}\big(\mathcal{R}_{[2]} \triangleright \mathcal{W}(g)\big) \otimes \mathcal{W}^{-1}\big(\mathcal{R}_{[1]} \triangleright \mathcal{W}(f)\big), \\
f(x^i) \odot_R g(y^j) &\equiv \mathcal{W}^{-1}\big(\mathcal{W}(g) \triangleleft \mathcal{R}_{[2]}\big) \otimes \mathcal{W}^{-1}\big(\mathcal{W}(f) \triangleleft \mathcal{R}_{[1]}\big),
\end{aligned} \tag{21}
$$

where f and g denote formal power series in the commutative coordinate algebras $\mathcal{A}_x$ and $\mathcal{A}_y$, respectively. The operations in (20) and (21) are referred to as *braided products.* Using the explicit form of the identities in (16) and (17) we derived in Ref. [42] commutation relations between normal ordered monomials. With these results at hand we are able to write down explicit formulae for computing braided products. As an example we give the expression we obtained for the braided product of the Manin plane [42]:

$$
\begin{aligned}
&f(x^1, x^2) \odot_L g(y^1, y^2) \\
&= \sum_{i=0}^{\infty} q^{i^2} (-\lambda)^i \, \frac{(y^2)^i \otimes (x^1)^i}{[[i]]_{q^{-2}}!} \, q^{-\hat{n}_{y^1} \otimes \hat{n}_{x^2} - 2\hat{n}_{y^2} \otimes \hat{n}_{x^1} - 2\hat{n}_{y^1} \otimes \hat{n}_{x^1} - 2\hat{n}_{y^2} \otimes \hat{n}_{x^1}}
\end{aligned}
$$

$$\times\,(D^1_{q^{-2}})^i g(q^{-i}y^1, q^{-2i}y^2)\otimes(D^2_{q^{-2}})^i f(q^{-2i}x^1, q^{-i}x^2). \tag{22}$$

In the above formulae we introduced the so-called *Jackson derivatives*, which are defined by [55]

$$D^i_{q^a} f(x^i) = \frac{f(x^i) - f(q^a x^i)}{(1-q^a)x^i}, \quad a\in\mathbb{C}. \tag{23}$$

It should be mentioned that the equalities in (12) and (13) imply that

$$\begin{aligned} f(x^i)\ \odot_L\ g(y^j) &= f(x^i)\ \odot_{\bar{R}}\ g(y^j),\\ f(x^i)\ \odot_{\bar{L}}\ g(y^j) &= f(x^i)\ \odot_R\ g(y^j). \end{aligned} \tag{24}$$

Thus, it would be sufficient to deal with $\odot_L$ and $\odot_{\bar{L}}$, only.

Now, we are in a position to introduce the tensor product of quantum spaces. It is equipped with a multiplication being determined by

$$(a\otimes a')(b\otimes b') = \big(a(\mathcal{R}_{[2]}\triangleright b)\big)\otimes\big((\mathcal{R}_{[1]}\triangleright a')b'\big), \tag{25}$$

or

$$(a\otimes a')(b\otimes b') = \big(a(\mathcal{R}^{-1}_{[1]}\triangleright b)\big)\otimes\big((\mathcal{R}^{-1}_{[2]}\triangleright a')b'\big), \tag{26}$$

where $a,b\in\mathcal{A}_q$ and $a',b'\in\mathcal{A}'_q$. We see that multiplication on a tensor product requires to know the commutation relations between the elements of the two tensor factors. Essentially for us is the fact that the algebra isomorphism $\mathcal{W}$ allows us to represent the tensor product of quantum spaces on a tensor product of commutative coordinate algebras. To be more specific, this can be achieved by extending the braided products in (20) and (21) as follows:

$$\begin{aligned} &\big(f(x^i)\otimes f'(y^j)\big)\ \odot_L\ \big(g(x^k)\otimes g'(y^l)\big)\\ &\equiv\big[f\overset{x}{\circledast}\mathcal{W}^{-1}\big(\mathcal{R}^{-1}_{[1]}\triangleright\mathcal{W}(g)\big)\big]\otimes\big[\mathcal{W}^{-1}\big(\mathcal{R}^{-1}_{[2]}\triangleright\mathcal{W}(f')\big)\overset{y}{\circledast}g'\big], \end{aligned} \tag{27}$$

and similarly

$$\begin{aligned} &\big(f(x^i)\otimes f'(y^j)\big)\ \odot_{\bar{L}}\ \big(g(x^k)\otimes g'(y^l)\big)\\ &\equiv\big[f\overset{x}{\circledast}\mathcal{W}^{-1}\big(\mathcal{R}_{[2]}\triangleright\mathcal{W}(g)\big)\big]\otimes\big[\mathcal{W}^{-1}\big(\mathcal{R}_{[1]}\triangleright\mathcal{W}(f')\big)\overset{y}{\circledast}g'\big], \end{aligned} \tag{28}$$

where $f, g \in \mathcal{A}_x$ and $f', g' \in \mathcal{A}_y$.

2.2 Definition and basic properties of q-deformed translations

Before introducing q-translations, it is useful to make contact with the notion of a *cross product algebra* [56]. It is well-known that we can combine a Hopf algebra $\mathcal{H}$ with its representation space $\mathcal{A}_q$ to form a left cross product algebra $\mathcal{A}_q \rtimes \mathcal{H}$ built on $\mathcal{A}_q \otimes \mathcal{H}$ with product

$$(a \otimes h)(b \otimes g) = a(h_{(1)} \triangleright b) \otimes h_{(2)}g, \quad a, b \in \mathcal{A}_q, \quad h, g \in \mathcal{H}. \tag{29}$$

There is also a right-handed version of this notion called a right cross product algebra $\mathcal{H} \ltimes \mathcal{A}$ and built on $\mathcal{H} \otimes \mathcal{A}$ with product

$$(h \otimes a)(g \otimes b) = hg_{(1)} \otimes (a \triangleleft g_{(2)})b. \tag{30}$$

When $\mathcal{A}_q$ is a q-deformed quantum space the cross product algebras have a Hopf structure. On quantum space generators the corresponding coproduct, antipode, and counit take the form [22, 23]

$$\begin{aligned} \Delta_{\bar{L}}(X^i) &= X^i \otimes 1 + (\bar{\mathcal{L}}_x)^i_j \otimes X^j, \\ \Delta_L(X^i) &= X^i \otimes 1 + (\mathcal{L}_x)^i_j \otimes X^j, \end{aligned} \tag{31}$$

$$\begin{aligned} S_{\bar{L}}(X^i) &= -S(\bar{\mathcal{L}}_x)^i_j X^j, \\ S_L(X^i) &= -S(\mathcal{L}_x)^i_j X^j, \end{aligned} \tag{32}$$

$$\varepsilon_{\bar{L}}(X^i) = \varepsilon_L(X^i) = 0. \tag{33}$$

There are opposite Hopf structures related to the above ones via

$$\Delta_{\bar{R}/R} = \tau \circ \Delta_{\bar{L}/L}, \qquad S_{\bar{R}/R} = S^{-1}_{\bar{L}/L}, \qquad \varepsilon_{\bar{R}/R} = \varepsilon_{\bar{L}/L}, \tag{34}$$

where τ shall denote the usual transposition of tensor factors. Notice that for the antipodes S_R and $S_{\bar{R}}$ it holds

$$\begin{aligned} S_{\bar{R}}(X^i) &= -X^j S^{-1}(\bar{\mathcal{L}}_x)^i_j, \\ S_R(X^i) &= -X^j S^{-1}(\mathcal{L}_x)^i_j, \end{aligned} \tag{35}$$

which is a direct consequence of the Hopf algebra axiom

$$a_{(2)}S^{-1}(a_{(1)}) = \varepsilon(a). \tag{36}$$

An essential observation is that coproducts of coordinates imply their translations [23,41,44,46,57,58]. This can be seen as follows. The coproduct on coordinates is an algebra homomorphism. If the coordinates constitute a module coalgebra then the algebra structure of the coordinates X^i is carried over to their coproduct $\Delta_A(X^i)$. More formally, we have

$$\Delta_A(X^iX^j) = \Delta_A(X^i)\Delta_A(X^j) \quad \text{and} \quad \Delta_A(h \triangleright X^i) = \Delta(h) \triangleright \Delta_A(X^i). \tag{37}$$

Due to this fact we can think of (31) as nothing other than an addition law for q-deformed vector components.

To proceed any further we have to realize that our algebra morphism $\mathcal{W}^{-1}$ can be extended by

$$\begin{gathered} \mathcal{W}_L^{-1} : \mathcal{A}_q \rtimes \mathcal{H} \longrightarrow \mathcal{A}, \\ \mathcal{W}_L^{-1}((X^1)^{i_1} \dots (X^n)^{i_n} \otimes h) \equiv \mathcal{W}^{-1}((X^1)^{i_1} \dots (X^n)^{i_n})\, \varepsilon(h), \end{gathered} \tag{38}$$

or

$$\begin{gathered} \mathcal{W}_R^{-1} : \mathcal{H} \ltimes \mathcal{A}_q \longrightarrow \mathcal{A}, \\ \mathcal{W}_R^{-1}(h \otimes (X^1)^{i_1} \dots (X^n)^{i_n}) \equiv \varepsilon(h)\, \mathcal{W}^{-1}((X^1)^{i_1} \dots (X^n)^{i_n}), \end{gathered} \tag{39}$$

with ε being the counit of the Hopf algebra $\mathcal{H}$. The meaning of the definitions in (38) and (39) becomes more clear when we assume the existence of a vacuum state $|0\rangle$ with

$$\begin{gathered} \langle 0|(X^1)^{i_1} \dots (X^n)^{i_n}|0\rangle = \mathcal{W}^{-1}((X^1)^{i_1} \dots (X^n)^{i_n}), \\ \langle 0|0\rangle = 1. \end{gathered} \tag{40}$$

Furthermore, we require for the vacuum state to be invariant under symmetry operations, i.e.

$$h|0\rangle = \varepsilon(h)|0\rangle. \tag{41}$$

Combining (38) and (39) then yields

$$\langle 0|(X^1)^{i_1} \dots (X^n)^{i_n} h|0\rangle = \langle 0|(X^1)^{i_1} \dots (X^n)^{i_n}|0\rangle \varepsilon(h)$$

$$= \mathcal{W}^{-1}((X^1)^{i_1} \dots (X^n)^{i_n})\varepsilon(h) = \mathcal{W}_L^{-1}((X^1)^{i_1} \dots (X^n)^{i_n} \otimes h). \tag{42}$$

Similar considerations hold for $\mathcal{W}_R^{-1}$.

With these mappings at hand we are able to introduce q-deformed translations:

$$\begin{aligned} f(x^i \oplus_{L/\bar{L}} y^j) &\equiv ((\mathcal{W}_L^{-1} \otimes \mathcal{W}_L^{-1}) \circ \Delta_{L/\bar{L}})(\mathcal{W}(f)), \\ f(x^i \oplus_{R/\bar{R}} y^j) &\equiv ((\mathcal{W}_R^{-1} \otimes \mathcal{W}_R^{-1}) \circ \Delta_{R/\bar{R}})(\mathcal{W}(f)), \end{aligned} \tag{43}$$

$$\begin{aligned} f(\ominus_{L/\bar{L}} x^i) &\equiv (\mathcal{W}_R^{-1} \circ S_{L/\bar{L}})(\mathcal{W}(f)), \\ f(\ominus_{R/\bar{R}} x^i) &\equiv (\mathcal{W}_L^{-1} \circ S_{R/\bar{R}})(\mathcal{W}(f)). \end{aligned} \tag{44}$$

The following formulae referring to the Manin plane shall serve as concrete examples:

$$f(x^i \oplus_L y^j) = \sum_{k_1,k_2=0}^{\infty} \frac{(x^1)^{k_1}(x^2)^{k_2}}{[[k_1]]_{q^{-2}}![[k_2]]_{q^{-2}}!} \left((D^1_{q^{-2}})^{k_1}(D^2_{q^{-2}})^{k_2} f\right)(q^{-k_2}y^1), \tag{45}$$

$$f(\ominus_L x^i) = q^{-(\hat{n}_{x^1})^2-(\hat{n}_{x^2})^2-2\hat{n}_{x^1}\hat{n}_{x^2}}\, f(-qx^1, -qx^2). \tag{46}$$

It should be mentioned that Eq. (45) can be seen as q-deformed version of the Taylor rule in two dimensions.

It is useful to notice that the considerations so far provide us with nothing other than a realization of a braided Hopf algebra on an algebra of commuting coordinates. For this reason we take another point of view which is provided by category theory. A category is a collection of objects $X, Y, Z, \dots$ together with a set $\mathrm{Mor}(X, Y)$ of morphisms between two objects X, Y. The composition of morphisms has similar properties as the composition of maps. We are interested in tensor categories. These categories have a product, denoted $\otimes$ and called the tensor product. It admits several 'natural' properties such as associativity and existence of a unit object. For a more formal treatment we refer to Refs. [33, 34, 47, 59]. If the action of a quasitriangular Hopf algebra $\mathcal{H}$ on the tensor product of two quantum spaces X and Y is defined by

$$h \triangleright (v \otimes w) = (h_{(1)} \triangleright v) \otimes (h_{(2)} \triangleright w) \in X \otimes Y, \quad h \in \mathcal{H}, \tag{47}$$

then the representations (quantum spaces) of the given Hopf algebra $\mathcal{H}$ are

the objects of a tensor category.

In this tensor category exist a number of morphisms of particular importance that are covariant with respect to the Hopf algebra action. First of all, for any pair of objects X, Y there is an isomorphism $\Psi_{X,Y} : X \otimes Y \to Y \otimes X$ such that $(g \otimes f) \circ \Psi_{X,Y} = \Psi_{X',Y'} \circ (f \otimes g)$ for arbitrary morphisms $f \in \mathrm{Mor}(X, X')$ and $g \in \mathrm{Mor}(Y, Y')$. In addition to this one requires the hexagon axiom to hold. The hexagon axiom is the validity of the two conditions

$$\Psi_{X,Z} \circ \Psi_{Y,Z} = \Psi_{X\otimes Y,Z}, \quad \Psi_{X,Z} \circ \Psi_{X,Y} = \Psi_{X,Y\otimes Z}. \tag{48}$$

A tensor category equipped with such mappings $\Psi_{X,Y}$ for each pair of objects X, Y is called a *braided tensor category*. The mappings $\Psi_{X,Y}$ as a whole are often referred to as the braiding of the tensor category. Furthermore, for any quantum space algebra X in this category there are morphisms $\Delta : X \to X \otimes X$, $S : X \to X$, and $\varepsilon : X \to \mathbb{C}$ forming a *braided Hopf algebra*, i.e. Δ, S, and ε obey the usual axioms of a Hopf algebra, but now as morphisms in the braided category. At this point, it should be noted that the inverse mappings $\Psi^{-1}_{X,Y}$ give a braiding as well, to which we can assign another braided Hopf structure denoted in the following by $\bar{\Delta} : X \to X \otimes X$, $\bar{S} : X \to X$, and $\bar{\varepsilon} : X \to \mathbb{C}$. For further details we recommend Refs. [47], [50], and [60].

In what follows it is important to realize that braided products and q-deformed translations are linked to the braiding mappings and the Hopf structures of a braided Hopf algebra, respectively. This way the axioms of a braided Hopf algebra carry over to the corresponding operations on commutative algebras. To this end, we first collect the axioms of a braided Hopf algebra, which are given by (see for example Ref. [35])

$$\begin{aligned} m \circ (S \otimes S) \circ \Psi^{-1} &= m \circ \Psi^{-1} \circ (S \otimes S) = S \circ m, \\ m \circ (\bar{S} \otimes \bar{S}) \circ \Psi &= m \circ \Psi \circ (\bar{S} \otimes \bar{S}) = \bar{S} \circ m, \end{aligned} \tag{49}$$

$$\begin{aligned} m \circ (S^{-1} \otimes S^{-1}) \circ \Psi &= m \circ \Psi \circ (S^{-1} \otimes S^{-1}) = S^{-1} \circ m, \\ m \circ (\bar{S}^{-1} \otimes \bar{S}^{-1}) \circ \Psi^{-1} &= m \circ \Psi^{-1} \circ (\bar{S}^{-1} \otimes \bar{S}^{-1}) = \bar{S}^{-1} \circ m, \end{aligned} \tag{50}$$

and

$$\begin{aligned} \Delta \circ S &= (S \otimes S) \circ (\Psi \circ \Delta), \\ \bar{\Delta} \circ \bar{S} &= (\bar{S} \otimes \bar{S}) \circ (\Psi^{-1} \circ \bar{\Delta}), \end{aligned} \tag{51}$$

$$
\begin{aligned}
(\Psi \circ \Delta) \circ S^{-1} &= (S^{-1} \otimes S^{-1}) \circ \Delta, \\
(\Psi^{-1} \circ \bar{\Delta}) \circ \bar{S}^{-1} &= (\bar{S}^{-1} \otimes \bar{S}^{-1}) \circ \bar{\Delta},
\end{aligned} \tag{52}
$$

where m denotes multiplication in the quantum space algebra. Furthermore, there are the axioms

$$
\begin{aligned}
m \circ (S \otimes \mathrm{id}) \circ \Delta = m \circ (\mathrm{id} \otimes S) \circ \Delta = \varepsilon, \\
m \circ (\bar{S} \otimes \mathrm{id}) \circ \bar{\Delta} = m \circ (\mathrm{id} \otimes \bar{S}) \circ \bar{\Delta} = \bar{\varepsilon},
\end{aligned} \tag{53}
$$

$$
\begin{aligned}
m \circ (S^{-1} \otimes \mathrm{id}) \circ (\Psi \circ \Delta) = m \circ (\mathrm{id} \otimes S^{-1}) \circ (\Psi \circ \Delta) = \varepsilon, \\
m \circ (\bar{S}^{-1} \otimes \mathrm{id}) \circ (\Psi^{-1} \circ \bar{\Delta}) = m \circ (\mathrm{id} \otimes \bar{S}^{-1}) \circ (\Psi^{-1} \circ \bar{\Delta}) = \bar{\varepsilon}.
\end{aligned} \tag{54}
$$

It is our aim to reformulate the above axioms in terms of the new operations on commutative coordinate algebras. For this to achieve we need a kind of dictionary which contains the correspondence between the set of morphisms in the braided category and the new operations on commutative algebras:

$$
\begin{aligned}
\Delta &\to f(x^i \oplus_L y^j), & S &\to f(\ominus_L x^i), \\
\bar{\Delta} &\to f(x^i \oplus_{\bar{L}} y^j), & \bar{S} &\to f(\ominus_{\bar{L}} x^i),
\end{aligned} \tag{55}
$$

$$
\begin{aligned}
\Psi \circ \Delta &\to f(x^i \oplus_R y^j), & S^{-1} &\to f(\ominus_R x^i), \\
\Psi^{-1} \circ \bar{\Delta} &\to f(x^i \oplus_{\bar{R}} y^j), & \bar{S}^{-1} &\to f(\ominus_{\bar{R}} x^i),
\end{aligned} \tag{56}
$$

$$
\begin{aligned}
\Psi &\to f(x^i) \odot_{\bar{L}} g(y^j), & \Psi^{-1} &\to f(x^i) \odot_L g(y^j), \\
m &\to f \circledast g.
\end{aligned} \tag{57}
$$

In addition to this, we should mention that

$$
\begin{aligned}
f(x^i \oplus_L y^j) = f(x^i \oplus_{\bar{R}} y^j), \\
f(x^i \oplus_{\bar{L}} y^j) = f(x^i \oplus_R y^j),
\end{aligned} \tag{58}
$$

and

$$
\begin{aligned}
f(\ominus_L x^i) = f(\ominus_{\bar{R}} x^i), \\
f(\ominus_{\bar{L}} x^i) = f(\ominus_R x^i).
\end{aligned} \tag{59}
$$

These identifications follow from arguments similar to those leading to (24) or

from a direct inspection of the explicit formulae for q-deformed translations [41].

Now, we are ready to rewrite the axioms in (49) and (50) as

$$\begin{aligned} f(\ominus_L\, x^i) \odot_L g(\ominus_L\, x^j) &= (f \circledast g)(\ominus_L\, x^i), \\ f(\ominus_{\bar{L}}\, x^i) \odot_{\bar{L}} g(\ominus_{\bar{L}}\, x^j) &= (f \circledast g)(\ominus_{\bar{L}}\, x^i), \end{aligned} \tag{60}$$

$$\begin{aligned} f(\ominus_R\, x^i) \odot_R g(\ominus_R\, x^j) &= (f \circledast g)(\ominus_R\, x^i), \\ f(\ominus_{\bar{R}}\, x^i) \odot_{\bar{R}} g(\ominus_{\bar{R}}\, x^j) &= (f \circledast g)(\ominus_{\bar{R}}\, x^i). \end{aligned} \tag{61}$$

Notice that we take the convention that tensor factors which are addressed by the same coordinates have to be multiplied via the star product. This means, for example,

$$f(x^i) \odot_{\bar{L}} g(x^j) = (\mathcal{R}_{[2]} \triangleright g(x^j)) \overset{x}{\circledast} (\mathcal{R}_{[1]} \triangleright f(x^i)). \tag{62}$$

Likewise we find from the axioms in (51) and (52) that

$$\begin{aligned} f(\ominus_L(x^i \oplus_L y^j)) &= f((\ominus_L\, x^i) \oplus_R (\ominus_L\, y^j)), \\ f(\ominus_{\bar{L}}(x^i \oplus_{\bar{L}} y^j)) &= f((\ominus_{\bar{L}}\, x^i) \oplus_{\bar{R}} (\ominus_{\bar{L}}\, y^j)), \end{aligned} \tag{63}$$

$$\begin{aligned} f(\ominus_R(x^i \oplus_R y^j)) &= f((\ominus_R\, x^i) \oplus_L (\ominus_R\, y^j)), \\ f(\ominus_{\bar{R}}(x^i \oplus_{\bar{R}} y^j)) &= f((\ominus_{\bar{R}}\, x^i) \oplus_{\bar{L}} (\ominus_{\bar{R}}\, y^j)). \end{aligned} \tag{64}$$

Furthermore, we are able to translate the axioms in (53) and (54) into the identities

$$\begin{aligned} f((\ominus_L\, x^i) \oplus_L x^j) &= f(x^i \oplus_L (\ominus_L\, x^j)) = f(0), \\ f((\ominus_{\bar{L}}\, x^i) \oplus_{\bar{L}} x^j) &= f(x^i \oplus_{\bar{L}} (\ominus_{\bar{L}}\, x^j)) = f(0), \end{aligned} \tag{65}$$

$$\begin{aligned} f((\ominus_R\, x^i) \oplus_R x^j) &= f(x^i \oplus_R (\ominus_R\, x^j)) = f(0), \\ f((\ominus_{\bar{R}}\, x^i) \oplus_{\bar{R}} x^j) &= f(x^i \oplus_{\bar{R}} (\ominus_{\bar{R}}\, x^j)) = f(0), \end{aligned} \tag{66}$$

where

$$f(0) \equiv \varepsilon(\mathcal{W}(f)) = f(x^i)\big|_{x^i=0}\,. \tag{67}$$

Notice that a function being q-translated depends on two sets of coordinates. In the above formulae the tensor factors corresponding to these coordinates

again have to be multiplied via the star product. The expressions for star products given in the work of Ref. [37] can easily be adapted to this task. In the case of the quantum plane, for example, we have

$$f(x^i, x^j) = q^{-\hat{n}_{x^2}\hat{n}_{y^1}}\ f(x^i, y^i)\big|_{y^i \to x^i}\,. \tag{68}$$

Finally, the property of the coproduct to be an algebra homomorphism [cf. (37)] gives rise to

$$\begin{aligned}(f \circledast g)(x^i \oplus_L y^j) &= f(x^i \oplus_L y^j)\ \odot_L\ g(x^i \oplus_L y^j),\\ (f \circledast g)(x^i \oplus_{\bar{L}} y^j) &= f(x^i \oplus_{\bar{L}} y^j)\ \odot_{\bar{L}}\ g(x^i \oplus_{\bar{L}} y^j),\end{aligned} \tag{69}$$

$$\begin{aligned}(f \circledast g)(x^i \oplus_R y^j) &= f(x^i \oplus_R y^j)\ \odot_R\ g(x^i \oplus_R y^j),\\ (f \circledast g)(x^i \oplus_{\bar{R}} y^j) &= f(x^i \oplus_{\bar{R}} y^j)\ \odot_{\bar{R}}\ g(x^i \oplus_{\bar{R}} y^j).\end{aligned} \tag{70}$$

To understand the right-hand side of the above equalities recall that each function being translated lives in a tensor product of two coordinate algebras. So functions that undergo the same q-translation have to be multiplied by the q-deformed tensor product of (27) or (28).

As we know, classical translations form a group. Considering q-deformed translations we can detect reminiscences of the group axioms satisfied by classical translations. We wish to illustrate this observation by the following calculation:

$$\begin{aligned}f((x^i \oplus_L y^j) \oplus_L (\ominus_L y^k)) &= f(x^i \oplus_L (y^j \oplus_L (\ominus_L y^k)))\\ &= f(x^i \oplus_L 0) = f(x^i).\end{aligned} \tag{71}$$

The first equality means associativity of q-deformed translations and follows from coassociativity of comultiplication. The second equality results from the identities in (65), which concern the existence of inverse elements. The last equality, which is a consequence of the Hopf algebra axiom

$$(\mathrm{id} \otimes \varepsilon) \circ \Delta = (\varepsilon \otimes \mathrm{id}) \circ \Delta = \mathrm{id}, \tag{72}$$

tells us that we have an identity element.

2.3 Conjugation properties and crossing symmetries

Now, we would like to discuss the conjugation properties of our quantum spaces. First of all, let us recall that our quantum spaces are endowed with a mapping $a \to \bar{a}$ that makes them $*$-algebras, i.e.

$$\overline{\alpha a + \beta b} = \bar{\alpha}\bar{a} + \bar{\beta}\bar{b}, \quad \overline{\bar{a}} = a, \quad \overline{ab} = \bar{b}\bar{a}, \tag{73}$$

with $\alpha, \beta \in \mathbb{C}$ and $a, b \in \mathcal{A}_q$. This mapping can be extended to tensor products of quantum spaces by

$$\overline{(a \otimes b)} = \bar{b} \otimes \bar{a}. \tag{74}$$

A quantum space generator is said to be real if

$$\overline{X^i} = X_i = g_{ij} X^j = \bar{X}_i, \tag{75}$$

where g_{ij} stands for the corresponding quantum metric. Notice that the quantum metric allows us to raise and lower indices as usual (see also Appendix A). From Eq. (75) we see that conjugating coordinates changes their transformation properties, i.e. contravariant coordinates become covariant and vice versa. However, there is one exception from (75) since the time element X^0 for q-deformed Minkowski space is imaginary, i.e. it fulfills

$$\overline{X^0} = -X_0 = -g_{00} X^0 = X^0 = -\bar{X}_0. \tag{76}$$

So $\mathrm{i}X^0$ is a real coordinate. In order to simplify our considerations we will assume that all quantum space coordinates are real. For q-deformed Minkowski space this requires to deal with $\mathrm{i}X^0$ instead of X^0.

Next, we wish to focus our attention on the conjugation properties of the L-matrices. Applying the conjugation properties of the symmetry generators to the explicit form of the L-matrices (see for example Ref. [43]) one can check the identities

$$\begin{aligned} \overline{(\mathcal{L}_x)^i_j} &= g_{ik}\, (\bar{\mathcal{L}}_x)^k_l\, g^{lj}, \\ \overline{(\bar{\mathcal{L}}_x)^i_j} &= g_{ik}\, (\mathcal{L}_x)^k_l\, g^{lj}. \end{aligned} \tag{77}$$

With these relations at hand we can proceed as follows:

$$\begin{aligned}\overline{X^i a} &= \overline{(1\otimes X^i)(a\otimes 1)} = \overline{\big((\bar{\mathcal{L}}_x)^i_j \triangleright a\big)\otimes X^j}\\ &= \bar{X}_j \otimes \big(\bar{a}\triangleleft \overline{(\bar{\mathcal{L}}_x)^i_j}\big) = \bar{X}_j \otimes \big(\bar{a}\triangleleft (\mathcal{L}_x)^k_l\, g_{ik} g^{lj}\big)\\ &= g_{ik}\,\bar{X}^l \otimes \big(\bar{a}\triangleleft (\mathcal{L}_x)^k_l\big) = g_{ik}(1\otimes\bar{a})(\bar{X}^k\otimes 1)\\ &= (1\otimes\bar{a})(\bar{X}_i\otimes 1) = \bar{a}\,\bar{X}_i. \end{aligned} \tag{78}$$

For arbitrary quantum space elements we analogously get

$$\begin{aligned}\overline{ab} &= \overline{(1\otimes a)(b\otimes 1)} = \overline{(\mathcal{R}_{[2]}\triangleright b)\otimes(\mathcal{R}_{[1]}\triangleright a)}\\ &= (\bar{a}\triangleleft\mathcal{R}_{[2]})\otimes(\bar{b}\triangleleft\mathcal{R}_{[1]}) = (1\otimes\bar{b})(\bar{a}\otimes 1)\\ &= \bar{b}\bar{a}, \end{aligned} \tag{79}$$

where for the third equality we used that the universal R-matrix is real in the sense

$$\overline{\mathcal{R}_{[1]}}\otimes\overline{\mathcal{R}_{[2]}} = \mathcal{R}_{[2]}\otimes\mathcal{R}_{[1]}. \tag{80}$$

It should be mentioned that the result in (79) remains unchanged if we use the braiding induced by $\tau\circ\mathcal{R}^{-1}$. The product in (78) and (79) should not be confused with that in (73). In (73) a and b live in the same quantum space, but in (78) and (79) this is not the case. Introducing the braiding mappings

$$\begin{aligned}\Psi_{\mathcal{A}_q,\mathcal{A}'_q}(a\otimes b) &\equiv (\mathcal{R}_{[2]}\triangleright b)\otimes(\mathcal{R}_{[1]}\triangleright a),\\ \Psi^{-1}_{\mathcal{A}_q,\mathcal{A}'_q}(a\otimes b) &\equiv (\mathcal{R}^{-1}_{[1]}\triangleright b)\otimes(\mathcal{R}^{-1}_{[2]}\triangleright a),\end{aligned} \tag{81}$$

the results of the above considerations can be reformulated as

$$\begin{aligned}\overline{\Psi_{\mathcal{A}_q,\mathcal{A}'_q}(a\otimes b)} &= \Psi_{\mathcal{A}_q,\mathcal{A}'_q}(\bar{b}\otimes\bar{a}),\\ \overline{\Psi^{-1}_{\mathcal{A}_q,\mathcal{A}'_q}(a\otimes b)} &= \Psi^{-1}_{\mathcal{A}_q,\mathcal{A}'_q}(\bar{b}\otimes\bar{a}).\end{aligned} \tag{82}$$

In very much the same way as in (78) we can compute for the coproduct of single quantum space generators

$$\begin{aligned}\overline{\Delta_L(X^i)} &= \overline{(X^i\otimes 1 + (\mathcal{L}_x)^i_j\otimes X^j)}\\ &= 1\otimes\bar{X}_i + \bar{X}_j\otimes\overline{(\mathcal{L}_x)^i_j}\\ &= 1\otimes\bar{X}_i + \bar{X}_j\otimes g_{ik}\,(\bar{\mathcal{L}}_x)^k_l\, g^{lj}\end{aligned}$$

$$= 1 \otimes g_{ik}\bar{X}^k + \bar{X}^l \otimes g_{ik}\,(\bar{\mathcal{L}}_x)^k_l$$
$$= \Delta_{\bar{R}}(g_{ik}\bar{X}^k) = \Delta_{\bar{R}}(\bar{X}_i). \tag{83}$$

This result confirms (cf. Ref. [50]),

$$\tau \circ (* \otimes *) \circ \Delta_{L/\bar{L}} = \Delta_{\bar{R}/R} \circ *, \tag{84}$$

where $*$ symbolizes the operation for conjugation. Similar considerations hold for the corresponding antipodes. As an example we give

$$\begin{aligned}\overline{S_L(X^i)} &= -\overline{S(\mathcal{L}_x)^i_j\, X^j} = -\bar{X}_j\, \overline{S(\mathcal{L}_x)^i_j} \\ &= -\bar{X}_j\, S^{-1}\overline{(\mathcal{L}_x)^i_j} = -\bar{X}_j\, g_{ik}\, g^{lj}\, S^{-1}(\bar{\mathcal{L}}_x)^k_l \\ &= -g_{ik}\,\bar{X}^l\, S^{-1}(\bar{\mathcal{L}}_x)^k_l = S_{\bar{R}}(\bar{X}_i),\end{aligned} \tag{85}$$

where the third equality makes use of the identity $* \circ S = S^{-1} \circ *$. The calculation in (85) and its counterparts for $S_{\bar{L}}$ and S_R imply

$$* \circ S_{L/\bar{L}} = S_{\bar{R}/R} \circ *. \tag{86}$$

Essentially for us is the fact that the conjugation on a quantum space $\mathcal{A}_q$ induces a mapping on the corresponding commutative algebra $\mathcal{A}$. This can be achieved through

$$\mathcal{W}(\,\overline{f(x^i)}\,) = \overline{\mathcal{W}(f(x^i))} \qquad \text{or} \qquad \overline{f(x^i)} = \mathcal{W}^{-1}(\,\overline{\mathcal{W}(f(x^i))}\,), \tag{87}$$

i.e. the algebra isomorphism $\mathcal{W}$ intertwines the conjugation on the quantum space algebra with that on the corresponding commutative algebra. Since we assume that a commutative function $f(x^i)$ can be written as a power series we especially have

$$\overline{f(x^i)} = \sum_{k_1,\ldots,k_n} \bar{f}_{k_1,\ldots,k_n}\, \overline{(x^1)^{k_1}\ldots(x^n)^{k_n}}, \tag{88}$$

where $\bar{f}_{k_1,\ldots,k_n}$ denotes the complex conjugate of the coefficient $f_{k_1,\ldots,k_n}$. Thus, it suffices to know how monomials behave under conjugation:

$$\begin{aligned}\overline{(x^1)^{k_1}\ldots(x^n)^{k_n}} &= \mathcal{W}^{-1}\big(\,\overline{(X^1)^{k_1}\ldots(X^n)^{k_n}}\,\big) \\ &= \mathcal{W}^{-1}\big(\,\overline{(X^n)^{k_n}}\ldots\overline{(X^1)^{k_1}}\,\big)\end{aligned}$$

$$
\begin{aligned}
&= \mathcal{W}^{-1}\big((X_n)^{k_n} \dots (X_1)^{k_1} \big) \\
&= \mathcal{W}^{-1}\big((g_{n\bar{n}} X^{\bar{n}})^{k_n} \dots (g_{1\bar{1}} X^{\bar{1}})^{k_1} \big) \\
&= \mathcal{W}^{-1}\big((g_{\bar{1}1} X^{1})^{k_n} \dots (g_{\bar{n}n} X^{n})^{k_1} \big) \\
&= (g_{\bar{1}1} x^{1})^{k_n} \dots (g_{\bar{n}n} x^{n})^{k_1}. \qquad (89)
\end{aligned}
$$

Notice that in the third equality we have introduced for each index i a so-called conjugate index $\overline{i}(\equiv n+1-i)$. For the quantum spaces we are considering for physical reasons the conjugate indices are given by

(i) (quantum plane) $\overline{i} \equiv 3-i, \quad i=1,2,$

(ii) (three-dimensional Euclidean space) $\overline{(+,3,-)} = (-,3,+),$

(iii) (four-dimensional Euclidean space) $\overline{i} \equiv 5-i, \quad i=1,\dots,4,$

(iv) (q-deformed Minkowski space) $\overline{(+,3,0,-)} = (-,3,0,+).$

One should also realize that there is no summation over repeated indices in (89), since the non-vanishing entries of the quantum metric are given by $g_{i\bar{i}}$ or $g_{\bar{i}i}$ for all possible values of i. Finally, we can summarize our considerations by

$$
\overline{f(x^i)} = \bar{f}(\,\overline{x^i}\,) \equiv \bar{f}(x^i)\big|_{x^i \to g_{i\bar{i}}\, x^{\bar{i}}}, \qquad (90)
$$

where

$$
\bar{f}(x^i) \sum_{k_1,\dots,k_n} \bar{f}_{k_1,\dots,k_n} (x^1)^{k_1} \dots (x^n)^{k_n}. \qquad (91)
$$

Now, we are in a position to describe the conjugation properties of star products, braided products, and q-deformed translations. First of all, from (73) and (87) it follows directly that

$$
\begin{aligned}
\overline{f(x^i) \circledast g(x^j)} &= \overline{\mathcal{W}^{-1}\big(\mathcal{W}(f)\,\mathcal{W}(g)\big)} = \mathcal{W}^{-1}\big(\overline{\mathcal{W}(f)\,\mathcal{W}(g)}\big) \\
&= \mathcal{W}^{-1}\big(\overline{\mathcal{W}(g)\,\mathcal{W}(f)}\big) = \mathcal{W}^{-1}\big(\mathcal{W}(\bar{g})\,\mathcal{W}(\bar{f})\big) \\
&= \mathcal{W}^{-1}\big(\mathcal{W}(\bar{g})\big)\mathcal{W}^{-1}\big(\mathcal{W}(\bar{f})\big) \\
&= \overline{g(x^i)} \circledast \overline{f(x^j)}. \qquad (92)
\end{aligned}
$$

It should also be clear from what we have done so far that the relations in (82) give rise to the following identities:

$$
\overline{f(x^i) \odot_L g(y^j)} = \overline{g(y^j)} \odot_{\bar{R}} \overline{f(x^i)} = \overline{g(y^j)} \odot_L \overline{f(x^i)},
$$

$$\overline{f(x^i)\odot_{\bar{L}}\ g(y^j)} = \overline{g(y^j)}\odot_R\ \overline{f(x^i)} = \overline{g(y^j)}\odot_{\bar{L}}\ \overline{f(x^i)}. \tag{93}$$

Furthermore, we can write down commutative counterparts of the relations in (84) and (86). Towards this end, we perform the calculation

$$\begin{aligned}
\overline{f(x^i \oplus_{L/\bar{L}} y^j)} &\overset{(43)}{=} \overline{\big(\mathcal{W}_L^{-1}\otimes\mathcal{W}_L^{-1}\big)\circ\Delta_{L/\bar{L}}(\mathcal{W}(f))} \\
&= \big(\mathcal{W}_R^{-1}\otimes\mathcal{W}_R^{-1}\big)\big(\overline{\Delta_{L/\bar{L}}(\mathcal{W}(f))}\big) \\
&\overset{(84)}{=} \big(\mathcal{W}_R^{-1}\otimes\mathcal{W}_R^{-1}\big)\big(\Delta_{\bar{R}/R}(\overline{\mathcal{W}(f)})\big) \\
&\overset{(87)}{=} \big(\mathcal{W}_R^{-1}\otimes\mathcal{W}_R^{-1}\big)\big(\Delta_{\bar{R}/R}(\mathcal{W}(\bar{f}))\big) \\
&= \bar{f}((\overline{y^j})\oplus_{\bar{R}/R}(\overline{x^i})),
\end{aligned} \tag{94}$$

and similarly

$$\begin{aligned}
\overline{f(\ominus_{L/\bar{L}}\, x^i)} &\overset{(44)}{=} \overline{\mathcal{W}_R^{-1}\big(S_{L/\bar{L}}(\mathcal{W}(f))\big)} = \mathcal{W}_L^{-1}\big(\overline{S_{L/\bar{L}}(\mathcal{W}(f))}\big) \\
&\overset{(86)}{=} \mathcal{W}_L^{-1}\big(S_{\bar{R}/R}(\overline{\mathcal{W}(f)})\big) \overset{(87)}{=} \mathcal{W}_L^{-1}\big(S_{\bar{R}/R}(\mathcal{W}(\bar{f}))\big) \\
&= \bar{f}(\ominus_{\bar{R}/R}(\overline{x^i})).
\end{aligned} \tag{95}$$

Notice that in (94) and (95) we make use of

$$\begin{aligned}
\overline{\mathcal{W}_L^{-1}(ah)} &= \mathcal{W}_R^{-1}(\bar{h}\bar{a}), \\
\overline{\mathcal{W}_R^{-1}(ha)} &= \mathcal{W}_L^{-1}(\bar{a}\bar{h}),
\end{aligned} \tag{96}$$

where $a\in\mathcal{A}_q$ and $h\in\mathcal{H}$. To sum up, the above considerations leave us with the rules

$$\begin{aligned}
\overline{f(x^i\oplus_{L/\bar{L}}\, y^j)} &= \bar{f}((\overline{y^j})\oplus_{\bar{R}/R}(\overline{x^i})), \\
\overline{f(x^i\oplus_{R/\bar{R}}\, y^j)} &= \bar{f}((\overline{y^j})\oplus_{\bar{L}/L}(\overline{x^i})),
\end{aligned} \tag{97}$$

$$\begin{aligned}
\overline{f(\ominus_{L/\bar{L}}\, x^i)} &= \bar{f}(\ominus_{\bar{R}/R}(\overline{x^i})), \\
\overline{f(\ominus_{R/\bar{R}}\, x^i)} &= \bar{f}(\ominus_{L/\bar{L}}(\overline{x^i})).
\end{aligned} \tag{98}$$

Our last comment in this section concerns the existence of crossing symmetries. They are based on the observation that the R-matrix and its inverse

are related to each other by

$$\hat{R}^{ij}_{kl} \overset{q\to 1/q}{\longleftrightarrow} (\hat{R}^{-1})^{\overline{i}\,\overline{j}}_{\overline{k}\,\overline{l}}. \tag{99}$$

This means that substituting $1/q$ for q we obtain from the entries of the R-matrix those of its inverse, but now labeled by the corresponding conjugate indices. The crossing symmetries are responsible for a number of remarkable correspondences. In the following we would like to present some of these correspondences.

In Ref. [37] we derived operator expressions for calculating star products. These formulae refer to a given normal ordering, but due to the crossing symmetries they can easily be transformed into expressions holding for reversed normal ordering. In other words, if we use as algebra homomorphism

$$\begin{aligned} \widetilde{\mathcal{W}} &: \mathcal{A} \longrightarrow \mathcal{A}_q, \\ \widetilde{\mathcal{W}}((x^1)^{i_1} \dots (x^n)^{i_n}) &= (X^n)^{i_n} \dots (X^1)^{i_1}, \end{aligned} \tag{100}$$

we get formulae for star products that can be obtained from those for the algebra homomorphism $\mathcal{W}$ [cf. (5)] via the transition

$$f \circledast g \overset{\substack{i \,\leftrightarrow\, \overline{i} \\ q \,\leftrightarrow\, 1/q}}{\longleftrightarrow} f \,\widetilde{\circledast}\, g, \tag{101}$$

where the symbol $\overset{\substack{i \,\leftrightarrow\, \overline{i} \\ q \,\leftrightarrow\, 1/q}}{\longleftrightarrow}$ now denotes the substitutions

$$D^i_{q^a} \to D^{\overline{i}}_{q^{-a}}, \quad \hat{n}_{x^i} \to \hat{n}_{x^{\overline{i}}}, \quad q \to q^{-1}. \tag{102}$$

Notice that the tilde in Eq. (101) shall remind us of the fact that the star product on the left-hand side refers to the algebra homomorphism $\widetilde{\mathcal{W}}$. In the case of the Manin plane the relationship in Eq. (101), for example, becomes

$$\begin{aligned} f(x^i) \circledast g(x^j) &= \left[q^{-\hat{n}_{x^2}\hat{n}_{y^1}} f(x^i) g(y^j)\right]_{y\to x} \\ \overset{\substack{i \,\leftrightarrow\, \overline{i} \\ q \,\leftrightarrow\, 1/q}}{\longleftrightarrow} & \left[q^{\hat{n}_{x^1}\hat{n}_{y^2}} f(x^i) g(y^j)\right]_{y\to x} = f(x^i) \,\widetilde{\circledast}\, g(x^j). \end{aligned} \tag{103}$$

In Refs. [41] and [42] we calculated explicit formulae for q-translations and braided products. The crossing symmetries we could read off from our results are sketched in Fig. 1. Notice that expressions linked by an arrow




Figure 1: Crossing-symmetries for braided products and q-translations

can be transformed into each other. The transition $i \leftrightarrow \bar{i}$ means that we have to substitute for each coordinate index the conjugate one. In the same way, $q \leftrightarrow 1/q$ requires to interchange q and its inverse. The symbol $x \leftrightarrow y$ indicates that the variables corresponding to different tensor factors have to be interchanged. Finally, the label 'ordering is reversed' tells us that the corresponding expressions refer to different algebra isomorphisms (given by $\mathcal{W}$ and $\widetilde{\mathcal{W}}$). We wish to illustrate these descriptions by the following example:

$$
\begin{aligned}
& f(x^i \oplus_L y^j) = \sum_{k_1,k_2=0}^{\infty} \frac{(x^1)^{k_1}(x^2)^{k_2}}{[[k_1]]_{q^{-2}}![[k_2]]_{q^{-2}}!} \left((D^1_{q^{-2}})^{k_1}(D^2_{q^{-2}})^{k_2} f\right)(q^{-k_2}y^1) \\
\overset{\substack{i \leftrightarrow \bar{i} \\ q \leftrightarrow 1/q}}{\longleftrightarrow} \quad & f(x^i \oplus_{\bar{L}} y^j) = \sum_{k_1,k_2=0}^{\infty} \frac{(x^2)^{k_2}(x^1)^{k_1}}{[[k_1]]_{q^2}![[k_2]]_{q^2}!} \left((D^1_{q^2})^{k_1}(D^2_{q^2})^{k_2} f\right)(q^{k_1}y^2) \\
\overset{\substack{x \leftrightarrow y \\ q \leftrightarrow 1/q}}{\longleftrightarrow} \quad & f(x^i \oplus_{\bar{R}} y^j) = \sum_{k_1,k_2=0}^{\infty} \frac{(y^1)^{k_1}(y^2)^{k_2}}{[[k_1]]_{q^{-2}}![[k_2]]_{q^{-2}}!} \left((D^1_{q^{-2}})^{k_1}(D^2_{q^{-2}})^{k_2} f\right)(q^{-k_1}y^2).
\end{aligned}
\tag{104}
$$

One should also keep in mind that the expressions for braided products and q-deformed translations depend on the choice for the normal ordering. Sometimes it is convenient to work with different normal orderings at the same time. However, a physical theory should be formulated in terms of objects referring to the same normal ordering. Thus, we need operators that enable us to transform a commutative function to another one representing the same quantum space element but now for reversed ordering. For the quantum spaces we are interested in the explicit form of these operators can be found in the work of Ref. [38]. In the case of the Manin plane, for example, they are given by

$$\begin{aligned} \tilde{f}(x^2, x^1) &= \hat{U}(f(x^1, x^2)) = q^{\hat{n}_{x^1}\hat{n}_{x^2}}\, f(x^1, x^2), \\ f(x^1, x^2) &= \hat{U}^{-1}(\tilde{f}(x^2, x^1)) = q^{-\hat{n}_{x^1}\hat{n}_{x^2}}\, \tilde{f}(x^2, x^1). \end{aligned} \tag{105}$$

Notice that we take the convention that the underlying normal ordering is indicated by the order in which the arguments of commutative functions are arranged.

3 Partial derivatives on quantum spaces

There are q-deformed analogs of partial derivatives, which act upon the algebra of quantum space coordinates [19–21]. In our approach the role of q-deformed partial derivatives and coordinates is completely symmetrical, i.e. partial derivatives show the same algebraic properties as quantum space coordinates. In this manner, q-deformed partial derivatives span quantum space algebras that are isomorphic to the quantum spaces they act upon. Thus, we can conclude that the considerations of the previous section also hold for the quantum space algebras of partial derivatives. In the present section we focus attention on actions of partial derivatives.

3.1 Leibniz rules and representations of partial derivatives

In analogy to (12) and (13) there are two possibilities for commutation relations between partial derivatives and quantum space coordinates. However, the action of partial derivatives on quantum space coordinates requires to modify the commutation relations in (12) and (13) in such a way that they

take the form

$$\partial^i X^j = g^{ij} + k(\hat{R}^{-1})^{ij}_{kl} X^k \partial^l,$$
$$\hat{\partial}^i X^j = \bar{g}^{ij} + k^{-1} \hat{R}^{ij}_{kl} X^k \hat{\partial}^l, \tag{106}$$

$$X^i \partial^j = -g^{ij} + k(\hat{R}^{-1})^{ij}_{kl} \partial^k X^l,$$
$$X^i \hat{\partial}^j = -\bar{g}^{ij} + k^{-1} \hat{R}^{ij}_{kl} \hat{\partial}^k X^l, \tag{107}$$

where we have introduced a conjugate quantum metric $\bar{g}^{ij}$. One should notice that in the case of the quantum plane it holds $\bar{g}^{ij} = -g^{ij}$. Otherwise we have $g^{ij} = \bar{g}^{ij}$. The constant k is a power of q and takes on the values

(i) (quantum plane) $k = q^2$,

(ii) (three-dimensional Euclidean space) $k = 1$,

(iii) (four-dimensional Euclidean space) $k = q$,

(iv) (q-deformed Minkowski space) $k = q^{-2}$.

Let us say a few words about the relationship between the partial derivatives ∂^i and $\hat{\partial}^i$. In the former literature [18, 48] ∂^i and $\hat{\partial}^i$ are seen to be conjugate to each other. In our approach, however, the partial derivatives ∂^i and $\hat{\partial}^i$ differ from each other by a normalization factor, only. Concretely, we have

$$\hat{\partial}^i = (\alpha_0)^{-1} k \partial^i, \tag{108}$$

where α_0 denotes the eigenvalue to the one-dimensional subspace of the vector representation of the universal R-matrix. For the quantum spaces we are interested in, relation (108) becomes

(i) (quantum plane) $\hat{\partial}^i = q^3 \partial^i$,

(ii) (three-dimensional Euclidean space) $\hat{\partial}^A = q^6 \partial^A$,

(iii) (four-dimensional Euclidean space) $\hat{\partial}^i = q^4 \partial^i$,

(iv) (q-deformed Minkowski space) $\hat{\partial}^\mu = q^{-4} \partial^\mu$.

From the q-deformed Leibniz rules in (106) and (107) we can derive right and left actions of partial derivatives on quantum space elements. To this end, we repeatedly apply the Leibniz rules to the product of a partial derivative with a normal ordered monomial of coordinates, until we obtain an expression with all partial derivatives standing to the right of all quantum space coordinates, i.e.

$$\partial^i (X^1)^{k_1} \dots (X^n)^{k_n} = \big(\partial^i_{(1)} \triangleright (X^1)^{k_1} \dots (X^n)^{k_n}\big) \partial^i_{(2)}. \tag{109}$$

Taking the counit of all partial derivatives appearing on the right-hand side finally yields the left action of ∂^i, since we have

$$\big(\partial^i_{(1)} \triangleright (X^1)^{k_1} \dots (X^n)^{k_n}\big) \varepsilon(\partial^i_{(2)}) = \partial^i \triangleright (X^1)^{k_1} \dots (X^n)^{k_n}. \tag{110}$$

Right actions of partial derivatives can be calculated in a similar way if we start from a partial derivative standing to the right of a normal ordered monomial and commute it to the left of all quantum space coordinates. Instead of (109) and (110), we now have

$$(X^1)^{k_1} \dots (X^n)^{k_n} \partial^i = \partial^i_{(2)} \big((X^1)^{k_1} \dots (X^n)^{k_n} \triangleleft \partial^i_{(1)}\big) \tag{111}$$

and

$$\varepsilon(\partial^i_{(2)}) \big((X^1)^{k_1} \dots (X^n)^{k_n} \triangleleft \partial^i_{(1)}\big) = (X^1)^{k_1} \dots (X^n)^{k_n} \triangleleft \partial^i. \tag{112}$$

The algebra isomorphism $\mathcal{W}$ allows us to introduce q-deformed derivatives that act upon commutative functions. By means of the relations

$$\begin{aligned} &\mathcal{W}(\partial^i \triangleright f) = \partial^i \triangleright \mathcal{W}(f), \quad f \in \mathcal{A}, \\ &\mathcal{W}(f \triangleleft \partial^i) = \mathcal{W}(f) \triangleleft \partial^i, \end{aligned} \tag{113}$$

or

$$\begin{aligned} \partial^i \triangleright f &\equiv \mathcal{W}^{-1}\left(\partial^i \triangleright \mathcal{W}(f)\right), \\ f \triangleleft \partial^i &\equiv \mathcal{W}^{-1}\left(\mathcal{W}(f) \triangleleft \partial^i\right), \end{aligned} \tag{114}$$

the actions of partial derivatives on the quantum space algebra $\mathcal{A}_q$ carry over to the corresponding commutative algebra $\mathcal{A}$. It should be obvious that each

Leibniz rule in (106) and (107) leads to its own q-derivative:

$$\begin{aligned} \partial^i X^j &= g^{ij} + k(\hat{R}^{-1})^{ij}_{kl}\, X^k \partial^l \qquad \Rightarrow \qquad \partial^i \triangleright f, \\ \hat{\partial}^i X^j &= \bar{g}^{ij} + k^{-1} \hat{R}^{ij}_{kl}\, X^k \hat{\partial}^l \qquad \Rightarrow \qquad \hat{\partial}^i \,\bar{\triangleright}\, f, \end{aligned} \tag{115}$$

$$\begin{aligned} X^i \partial^j &= -\bar{g}^{ij} + k(\hat{R}^{-1})^{ij}_{kl}\, \partial^k X^l \qquad \Rightarrow \qquad f \,\bar{\triangleleft}\, \partial^i, \\ X^i \hat{\partial}^j &= -g^{ij} + k^{-1} \hat{R}^{ij}_{kl}\, \hat{\partial}^k X^l \qquad \Rightarrow \qquad f \triangleleft \hat{\partial}^i. \end{aligned} \tag{116}$$

In the work of Ref. [38] we derived operator representations for q-deformed partial derivatives by applying these ideas. Our results can be viewed as multi-dimensional versions of the celebrated Jackson derivative [55]. As an example we write down the left representations for partial derivatives on the two-dimensional quantum plane:

$$\begin{aligned} \partial^1 \triangleright f(x^1, x^2) &= -q^{-1/2} D^2_{q^2} f(qx^1, x^2), \\ \partial^2 \triangleright f(x^1, x^2) &= q^{1/2} D^1_{q^2} f(x^1, q^2 x^2). \end{aligned} \tag{117}$$

Next, we would like to discuss a point that is important for q-deformed quantum mechanics. In our approach, as already mentioned, there is a symmetry between coordinates and partial derivatives with the exception that under conjugation coordinates are real objects, while partial derivatives have to be imaginary (see the discussion in the next subsection). However, if we introduce momentum variables by setting $P^k \equiv \mathrm{i}\partial^k$, we obtain objects with the very same conjugation properties as quantum space coordinates. With these new variables the Leibniz rules in (106) and (107) now become

$$\begin{aligned} P^k(\mathrm{i}X^l) &= -g^{kl} + k(\hat{R}^{-1})^{kl}_{mn}\, (\mathrm{i}X^m) P^n, \\ P^k(\mathrm{i}\hat{X}^l) &= -\bar{g}^{kl} + k^{-1} \hat{R}^{kl}_{mn}\, (\mathrm{i}\hat{X}^m) P^n, \end{aligned} \tag{118}$$

$$\begin{aligned} (\mathrm{i}X^k) P^l &= \bar{g}^{kl} + k(\hat{R}^{-1})^{kl}_{mn}\, P^m (\mathrm{i}X^n), \\ (\mathrm{i}\hat{X}^k) P^l &= g^{kl} + k^{-1} \hat{R}^{kl}_{mn}\, P^m (\mathrm{i}\hat{X}^n). \end{aligned} \tag{119}$$

This means that

$$\partial^k_p \equiv \mathrm{i}X^k \quad \text{as well as} \quad \hat{\partial}^k_p \equiv \mathrm{i}\hat{X}^k \tag{120}$$

can play the role of partial derivatives on momentum space. In the case of the Manin plane, however, we have to take into account an additional minus

sign in the definition of ∂_p^k and $\hat{\partial}_p^k$.

Furthermore, we see by comparing (118) with (106) and (119) with (107) that the Leibniz rules for position and momentum space are linked via the substitutions

$$X^k \longleftrightarrow P^k, \quad \partial_x^k \longleftrightarrow \partial_p^k, \quad \hat{\partial}_x^k \longleftrightarrow \hat{\partial}_p^k. \tag{121}$$

Due to this fact, the actions of partial derivatives on momentum space take on the same form as those for partial derivatives on coordinate space. To be more specific we have the correspondences

$$\begin{aligned} \partial_x^k \triangleright f(x^j) &\overset{x\leftrightarrow p}{\longleftrightarrow} \partial_p^k \triangleright f(p^j), \\ \hat{\partial}_x^k \,\bar{\triangleright}\, f(x^j) &\overset{x\leftrightarrow p}{\longleftrightarrow} \hat{\partial}_p^k \,\bar{\triangleright}\, f(p^j), \end{aligned} \tag{122}$$

$$\begin{aligned} f(x^j) \triangleleft \hat{\partial}_x^k &\overset{x\leftrightarrow p}{\longleftrightarrow} f(p^j) \triangleleft \hat{\partial}_p^k, \\ f(x^j) \,\bar{\triangleleft}\, \partial_x^k &\overset{x\leftrightarrow p}{\longleftrightarrow} f(p^j) \,\bar{\triangleleft}\, \partial_p^k, \end{aligned} \tag{123}$$

where $\overset{x\leftrightarrow p}{\longleftrightarrow}$ instructs us to substitute the momentum variable p^i for each space coordinate x^i and vice versa. These observations should tell us that all considerations for position space pertain equally for momentum space.

3.2 Conjugation properties and crossing symmetries

As already mentioned, partial derivatives obey the same algebra relations as the corresponding quantum space coordinates. Since we require for partial derivatives to be imaginary, they have to satisfy

$$\overline{\partial^i} = -\bar{\partial}_i = -\partial_i = -g_{ij}\partial^j. \tag{124}$$

Again, there is one exception from this rule, since in the case of the q-deformed Minkowski space we have

$$\overline{\partial^0} = \bar{\partial}_0 = \partial_0 = g_{00}\partial^0 = -\partial^0. \tag{125}$$

Next, we would like to say a few words about the conjugation properties of the actions of partial derivatives. To this aim, we examine how the Leibniz rules in (106) and (107) change under conjugation. We have

$$\overline{\partial^i X^j} = \overline{g^{ij} + k(\hat{R}^{-1})^{ij}_{kl}\, X^k \partial^l}$$

$$
\begin{aligned}
\Rightarrow -\bar{X}_j\bar{\partial}_i &= \bar{g}_{ij} - k(\hat{R}^{-1})^{kl}_{ij}\,\bar{\partial}_l\bar{X}_k \\
\Rightarrow X_j\partial_i &= -\bar{g}_{ij} + k(\hat{R}^{-1})^{kl}_{ij}\,\partial_l X_k \\
\Rightarrow X^i\partial^j &= -\bar{g}^{ij} + k(\hat{R}^{-1})^{ij}_{kl}\,\partial^k X^l,
\end{aligned} \tag{126}
$$

and likewise

$$
\begin{aligned}
\overline{\hat{\partial}^i X^j} &= \overline{\bar{g}^{ij} + k^{-1}\hat{R}^{ij}_{kl}\,X^k\hat{\partial}^l} \\
\Rightarrow X^i\hat{\partial}^j &= -g^{ij} + k^{-1}\hat{R}^{ij}_{kl}\,\hat{\partial}^k X^l.
\end{aligned} \tag{127}
$$

The derivation in (126) makes use of (73), (75), and (124). In addition to this, we need the identities

$$
\begin{aligned}
\overline{g^{ij}} = g^{ij} = \bar{g}_{ij}, \quad \overline{\bar{g}^{ij}} = \bar{g}^{ij} = g_{ij}, \\
\overline{(\hat{R}^{\pm 1})^{kl}_{ij}} = (\hat{R}^{\pm 1})^{kl}_{ij} = (\hat{R}^{\pm 1})^{ij}_{kl}, \\
g^{ml}g^{nk}(\hat{R}^{\pm 1})^{ij}_{kl} = (\hat{R}^{\pm 1})^{mn}_{kl}\,g^{li}g^{kj}.
\end{aligned} \tag{128}
$$

The above considerations tell us that under conjugation the Leibniz rules leading to left actions transform into those for right actions and vice versa. However, this statement is only valid, if we deal with coordinates and partial derivatives being respectively real and imaginary under conjugation. For q-deformed Minkowski space this means that we have to work with $\mathrm{i}X^0$ and $\mathrm{i}\partial^0$ instead of X^0 and ∂^0. In doing so, the quantum metric and the R-matrices for q-deformed Minkowski space must be modified as follows:

$$
\begin{aligned}
g^{\mu\nu} &\to \mathrm{i}^{\delta_{\mu 0}+\delta_{\nu 0}}\,g^{\mu\nu}, \\
(\hat{R}^{\pm 1})^{\mu\nu}_{\rho\sigma} &\to \mathrm{i}^{\delta_{\mu 0}+\delta_{\nu 0}}(\hat{R}^{\pm 1})^{\mu\nu}_{\rho\sigma}\,\mathrm{i}^{\delta_{\rho 0}+\delta_{\sigma 0}}.
\end{aligned} \tag{129}
$$

The substitutions in (129) can be viewed as a q-analog of classical Wick rotation. By virtue of the identifications in (115) and (116), we finally conclude that the results of (126) and (127) imply the identities

$$
\begin{aligned}
\overline{\partial^i \triangleright f(x^j)} &= -\overline{f(x^j)}\,\bar{\triangleleft}\,\partial_i, & \overline{f(x^j)\,\bar{\triangleleft}\,\partial^i} &= -\partial_i \triangleright \overline{f(x^j)}, \\
\overline{\hat{\partial}^i\,\bar{\triangleright}\,f(x^j)} &= -\overline{f(x^j)}\,\triangleleft\,\hat{\partial}_i, & \overline{f(x^j)\,\triangleleft\,\hat{\partial}^i} &= -\hat{\partial}_i\,\bar{\triangleright}\,\overline{f(x^j)}.
\end{aligned} \tag{130}
$$

Next, we turn to the crossing symmetries for partial derivatives, since they give a simple method to transform the different representations of par-

tial derivatives into each other. In the case of partial derivatives crossing symmetries represent the fact that we can make a transition between a Leibniz rule and its conjugate version by applying the substitutions

$$\partial^i \leftrightarrow \hat{\partial}^{\overline{i}}, \quad X^i \leftrightarrow X^{\overline{i}}, \quad q \leftrightarrow q^{-1}. \tag{131}$$

For this to become more clear we write down the explicit form of the Leibniz rules in the case of the two-dimensional quantum plane:

$$\begin{aligned} \partial^1 X^1 &= q X^1 \partial^1, \\ \partial^1 X^2 &= -q^{-1/2} + q^2 X^2 \partial^1, \end{aligned} \tag{132}$$

$$\begin{aligned} \partial^2 X^1 &= q^{1/2} + q^2 X^1 \partial^2 - q^2 \lambda X^2 \partial^1, \\ \partial^2 X^2 &= q X^2 \partial^2. \end{aligned} \tag{133}$$

The substitutions in (131) indeed give the conjugate Leibniz rules, i.e.

$$\begin{aligned} \hat{\partial}^1 X^1 &= q^{-1} X^1 \hat{\partial}^1, \\ \hat{\partial}^1 X^2 &= q^{-1/2} + q^{-2} X^2 \hat{\partial}^1 + q^{-2} \lambda X^1 \hat{\partial}^2, \end{aligned} \tag{134}$$

$$\begin{aligned} \hat{\partial}^2 X^1 &= -q^{1/2} + q^{-2} X^1 \hat{\partial}^2, \\ \hat{\partial}^2 X^2 &= q^{-1} X^2 \hat{\partial}^2. \end{aligned} \tag{135}$$

This correspondence is the deeper reason for the transition rules:

$$\begin{aligned} & \partial^1 \triangleright f(x^1, x^2) = -q^{-1/2} D^2_{q^2} f(q x^1, x^2) \\ \overset{\substack{i \to \overline{i} \\ q \to 1/q}}{\longleftrightarrow} \quad & \hat{\partial}^2 \bar{\triangleright} f(x^2, x^1) = -q^{1/2} D^1_{q^{-2}} f(q^{-1} x^2, x^1), \end{aligned} \tag{136}$$

$$\begin{aligned} & \partial^2 \triangleright f(x^1, x^2) = q^{1/2} D^1_{q^2} f(x^1, q^2 x^2) \\ \overset{\substack{i \to \overline{i} \\ q \to 1/q}}{\longleftrightarrow} \quad & \hat{\partial}^1 \bar{\triangleright} f(x^2, x^1) = q^{-1/2} D^2_{q^{-2}} f(x^2, q^{-2} x^1), \end{aligned} \tag{137}$$

where the transition symbol has the same meaning as in (102). It should be noted that the above transition rules require to reverse the underlying normal ordering.

It is also possible to establish a relationship between right and left ac-

tions of partial derivatives. To this end, let us recall that the identities in (130) enable us to derive right representations from left ones and vice versa. Exploiting this idea one can show that in the case of the Manin plane we have

$$\begin{aligned} & \partial^1 \triangleright f(x^1, x^2) = -q^{-1/2} D^2_{q^2} f(qx^1, x^2) \\ \overset{i \leftrightarrow \bar{i}}{\longleftrightarrow} \quad & -f(x^1, x^2) \,\bar{\triangleleft}\, \partial^2 = q^{-1/2} D^1_{q^2} f(x^1, qx^2), \end{aligned} \tag{138}$$

$$\begin{aligned} & \partial^2 \triangleright f(x^1, x^2) = q^{1/2} D^1_{q^2} f(x^1, q^2 x^2) \\ \overset{i \leftrightarrow \bar{i}}{\longleftrightarrow} \quad & -f(x^1, x^2) \,\bar{\triangleleft}\, \partial^1 = -q^{1/2} D^2_{q^2} f(q^2 x^1, x^2), \end{aligned} \tag{139}$$

where the symbol $\overset{i \leftrightarrow \bar{i}}{\longleftrightarrow}$ stands for the substitutions

$$D^i_{q^a} \to D^{\bar{i}}_{q^a}, \quad \hat{n}_{x^i} \to \hat{n}_{x^{\bar{i}}}, \quad x^i \to x^{\bar{i}}. \tag{140}$$

These reasonings carry over to the other quantum spaces we are considering without any problem. In general, the correspondences between the different representations diagrammatically appear as in Fig. 2. For the sake of completeness we want to list the normal orderings that lead to the simplest expressions for unconjugate left actions of partial derivatives:

(i) (quantum plane) $(X^1)^{n_1}(X^2)^{n_2}$,

(ii) (three-dimensional Euclidean space) $(X^+)^{n_+}(X^3)^{n_3}(X^-)^{n_-}$,

(iii) (four-dimensional Euclidean space) $(X^4)^{n_4}(X^3)^{n_3}(X^2)^{n_2}(X^1)^{n_1}$,

(iv) (q-deformed Minkowski space) $(X^-)^{n_-}(X^{3/0})^{n_{3/0}}(X^3)^{n_3}(X^+)^{n_+}$.

3.3 Further correspondences between actions of partial derivatives

Next, we would like to mention that the operations introduced in (44) intertwine the actions of partial derivatives with their conjugate counterparts, as was pointed out in Ref. [50]. Using the explicit form for the actions of partial derivatives and that for the operations in (44) one directly verifies that

$$\hat{\partial}^i \,\bar{\triangleright}\, \big(f(\ominus_R\, x^j)\big) = -(\partial^i \triangleright f)(\ominus_R\, x^j),$$




Figure 2: Crossing-symmetries for q-deformed partial derivatives

$$\partial^i \triangleright \big(f(\ominus_{\bar{R}}\, x^j)\big) = -(\hat{\partial}^i \,\bar{\triangleright}\, f)(\ominus_{\bar{R}}\, x^j), \tag{141}$$

$$\begin{aligned} \big(f(\ominus_{\bar{L}}\, x^j)\big) \triangleleft \hat{\partial}^i &= -(f \,\bar{\triangleleft}\, \partial^i)(\ominus_{\bar{L}}\, x^j), \\ \big(f(\ominus_L\, x^j)\big) \,\bar{\triangleleft}\, \partial^i &= -(f \triangleleft \hat{\partial}^i)(\ominus_L\, x^j). \end{aligned} \tag{142}$$

It should be noticed that in the case of the Manin plane the minus sign on the right-hand side must be dropped.

From (141) and (142) we can derive further relations which also intertwine the actions of partial derivatives with their conjugate versions. To clarify this assertion we start from the first identity in (141) and proceed as follows:

$$\begin{aligned} & \hat{\partial}^i \,\bar{\triangleright}\, \big(f(\ominus_R\, x^j)\big) = -(\partial^i \triangleright f)(\ominus_R\, x^j), \\ \Rightarrow \quad & \hat{\partial}^i \,\bar{\triangleright}\, S^{-1}(\mathcal{W}(f)) = -S^{-1}(\partial^i \triangleright \mathcal{W}(f)), \\ \Rightarrow \quad & S\big(\hat{\partial}^i \,\bar{\triangleright}\, S^{-1}(\mathcal{W}(f))\big) = -\partial^i \triangleright \mathcal{W}(f), \\ \Rightarrow \quad & S\big(\hat{\partial}^i \,\bar{\triangleright}\, \mathcal{W}(f)\big) = -\partial^i \triangleright S(\mathcal{W}(f)), \\ \Rightarrow \quad & (\hat{\partial}^i \,\bar{\triangleright}\, f)(\ominus_L\, x^j) = -\partial^i \triangleright \big(f(\ominus_L\, x^j)\big). \end{aligned} \tag{143}$$

The first step uses (44) and (114). For the second step we hit both sides of the second relation by the antipode. For the third step we substitute $S(\mathcal{W}(f))$ for $\mathcal{W}(f)$ and the final result is obtained by applying the definitions in (114). Repeating the same steps for the other identities in (141) and (142) gives us

$$(\partial^i \triangleright f)(\ominus_{\bar{L}}\, x^j) = -\hat{\partial}^i \,\bar{\triangleright}\, \big(f(\ominus_{\bar{L}}\, x^j)\big), \tag{144}$$

$$\begin{aligned}(f \triangleleft \hat{\partial}^i)(\ominus_{\bar{R}}\, x^j) &= -\big(f(\ominus_{\bar{R}}\, x^j)\big) \,\bar{\triangleleft}\, \partial^i,\\ (f \,\bar{\triangleleft}\, \partial^i)(\ominus_{R}\, x^j) &= -\big(f(\ominus_{R}\, x^j)\big) \triangleleft \hat{\partial}^i.\end{aligned} \tag{145}$$

Our last comment in this section concerns the fact that we can switch between right and left representations of partial derivatives by means of the antipode and its inverse. Concretely, we have the identities

$$\begin{aligned}\partial^i \triangleright f(x^k) &= f(x^k) \triangleleft S_L(\partial^i) = -\big(f(x^k) \triangleleft S(\mathcal{L}_\partial)^i_j\big) \triangleleft \partial^j\\ &= -\big((\mathcal{L}_\partial)^i_j \triangleright f(x^k)\big) \triangleleft \partial^j,\\ \partial^i \,\bar{\triangleright}\, f(x^k) &= f(x^k) \,\bar{\triangleleft}\, S_{\bar{L}}(\partial^i) = -\big(f(x^k) \triangleleft S(\bar{\mathcal{L}}_\partial)^i_j\big) \,\bar{\triangleleft}\, \partial^j\\ &= -\big((\bar{\mathcal{L}}_\partial)^i_j \triangleright f(x^k)\big) \,\bar{\triangleleft}\, \partial^j,\end{aligned} \tag{146}$$

$$\begin{aligned}f(x^k) \triangleleft \partial^i &= S_R(\partial^i) \triangleright f(x^k) = -\partial^j \triangleright \big(S^{-1}(\mathcal{L}_\partial)^i_j \triangleright f(x^k)\big)\\ &= -\partial^j \triangleright \big(f(x^k) \triangleleft (\mathcal{L}_\partial)^i_j\big),\\ f(x^k) \,\bar{\triangleleft}\, \partial^i &= S_{\bar{R}}(\partial^i) \,\bar{\triangleright}\, f(x^k) = -\partial^j \,\bar{\triangleright}\, \big(S^{-1}(\bar{\mathcal{L}}_\partial)^i_j \triangleright f(x^k)\big)\\ &= -\partial^j \,\bar{\triangleright}\, \big(f(x^k) \triangleleft (\bar{\mathcal{L}}_\partial)^i_j\big),\end{aligned} \tag{147}$$

where we used (32) and (35).

4 Dual pairings and q-exponentials

In this section we deal with dual pairings and exponentials for q-deformed quantum spaces. Let us recall that the comodules of a given quantum group form the objects of a braided tensor category [cf. Section 2]. In what follows we suppose that for each quantum space algebra $\mathcal{A}_q$ in the braided tensor category we have a dual object $\mathcal{A}_q^*$ living in the same category. Our assumption

means that in our tensor category we have dual pairings

$$\begin{aligned} \langle .\,,. \rangle &: \mathcal{A}_q \otimes \mathcal{A}_q^* \to \mathbb{C} \quad \text{with} \quad \left\langle e_a, f^b \right\rangle = \delta_a^b, \\ \langle .\,,. \rangle' &: \mathcal{A}_q^* \otimes \mathcal{A}_q \to \mathbb{C} \quad \text{with} \quad \left\langle f^b, e_a \right\rangle' = \delta_a^b, \end{aligned} \tag{148}$$

where $\{e_a\}$ is a basis in $\mathcal{A}_q$ and $\{f^b\}$ a dual basis in $\mathcal{A}_q^*$.

4.1 Definition and basic properties of dual pairings

In Ref. [46] it was shown that the algebra of quantum space coordinates and that of the corresponding partial derivatives are dual to each other. The dual pairing is given by

$$\langle .\,,. \rangle : \mathcal{A}_q^* \otimes \mathcal{A}_q \to \mathbb{C} \quad \text{with} \quad \left\langle f(\partial^i), g(X^j) \right\rangle \equiv \varepsilon(f(\partial^i) \triangleright g(X^j)), \tag{149}$$

or

$$\langle .\,,. \rangle' : \mathcal{A}_q \otimes \mathcal{A}_q^* \to \mathbb{C} \quad \text{with} \quad \left\langle f(X^i), g(\partial^j) \right\rangle' \equiv \varepsilon(f(X^i) \triangleleft g(\partial^j)). \tag{150}$$

As usual the above pairings carry over to commutative algebras by means of the algebra isomorphism $\mathcal{W}$, i.e.

$$\langle .\,,. \rangle : \mathcal{A}^* \otimes \mathcal{A} \to \mathbb{C} \quad \text{with} \quad \left\langle f(\partial^i), g(x^j) \right\rangle \equiv \left\langle \mathcal{W}(f(\partial^i)), \mathcal{W}(g(x^j)) \right\rangle, \tag{151}$$

and likewise

$$\langle .\,,. \rangle' : \mathcal{A} \otimes \mathcal{A}^* \to \mathbb{C} \quad \text{with} \quad \left\langle f(x^i), g(\partial^j) \right\rangle' \equiv \left\langle \mathcal{W}(f(x^i)), \mathcal{W}(g(\partial^j)) \right\rangle'. \tag{152}$$

Since the partial derivatives can act on coordinates in four different ways, there are four possibilities for defining a pairing between coordinates and derivatives. More concretely, we have the pairings

$$\begin{aligned} \left\langle f(\partial^i), g(x^j) \right\rangle_{L,\bar{R}} &\equiv (f(\partial^i) \triangleright g(x^j))|_{x^j=0} = (f(\partial^i) \,\bar{\triangleleft}\, g(x^j))|_{\partial^i=0}, \\ \left\langle f(\hat{\partial}^i), g(x^j) \right\rangle_{\bar{L},R} &\equiv (f(\hat{\partial}^i) \,\bar{\triangleright}\, g(x^j))|_{x^j=0} = (f(\hat{\partial}^i) \triangleleft g(x^j))|_{\partial^i=0}, \end{aligned} \tag{153}$$

$$\begin{aligned} \left\langle f(x^i), g(\partial^j) \right\rangle_{L,\bar{R}} &\equiv (f(x^i) \,\bar{\triangleleft}\, g(\partial^j))|_{x^i=0} = (f(x^i) \triangleright g(\partial^j))|_{\partial^j=0}, \\ \left\langle f(x^i), g(\hat{\partial}^j) \right\rangle_{\bar{L},R} &\equiv (f(x^i) \triangleleft g(\hat{\partial}^j))|_{x^i=0} = (f(x^i) \,\bar{\triangleright}\, g(\hat{\partial}^j))|_{\partial^j=0}. \end{aligned} \tag{154}$$

Notice that in (153) and (154) a label is introduced to characterize the different pairings. Its meaning can be understood in the following way. The left and right indices refer to the left and right arguments of the dual parings, respectively. If an argument of a given dual paring acts on the other argument via the conjugate representation the index for the acting argument is overlined. One should also notice that each pairing can be calculated in two different ways. This observation shows us the symmetry between derivatives and coordinates once again. As an example we give the explicit form of a pairing for the Manin plane. In this case the first pairing in (153) reads on normal ordered monomials as follows:

$$\left\langle (\partial_1)^{n_1}(\partial_2)^{n_2}, (x^2)^{m_2}(x^1)^{m_1} \right\rangle_{L,\bar{R}} = \delta_{m_1,n_1}\delta_{m_2,n_2}\,[[n_1]]_{q^2}!\,[[n_2]]_{q^2}!. \tag{155}$$

From their definitions in (153) and (154) we see that the different pairings are determined by the actions of derivatives on coordinates. On the other hand the dual pairings together with the coaddition on quantum spaces allow us to write down the so-called coregular actions, which are identical with the actions in (115) and (116). Concretely, we have

$$\begin{aligned}
\partial^i \triangleright f(x^j) &= \left\langle \partial^i, f_{(\bar{L},1)} \right\rangle_{L,\bar{R}} f_{(\bar{L},2)} = \left(\partial^i \overset{y}{\triangleright} f(y^k \oplus_{\bar{L}} x^j) \right)\Big|_{y^k=0}, \\
\hat{\partial}^i \bar{\triangleright} f(x^j) &= \left\langle \hat{\partial}^i, f_{(L,1)} \right\rangle_{\bar{L},R} f_{(L,2)} = \left(\hat{\partial}^i \overset{y}{\bar{\triangleright}} f(y^k \oplus_{L} x^j) \right)\Big|_{y^k=0},
\end{aligned} \tag{156}$$

$$\begin{aligned}
f(x^j) \bar{\triangleleft} \partial^i &= f_{(R,1)} \left\langle f_{(R,2)}, \partial^i \right\rangle_{L,\bar{R}} = \left(f(x^j \oplus_R y^k) \overset{y}{\bar{\triangleleft}} \partial^i \right)\Big|_{y^k=0}, \\
f(x^j) \triangleleft \hat{\partial}^i &= f_{(\bar{R},1)} \left\langle f_{(\bar{R},2)}, \hat{\partial}^i \right\rangle_{\bar{L},R} = \left(f(x^j \oplus_{\bar{R}} y^k) \overset{y}{\triangleleft} \hat{\partial}^i \right)\Big|_{y^k=0},
\end{aligned} \tag{157}$$

where

$$\begin{aligned}
f_{(L/\bar{L},1)} \otimes f_{(L/\bar{L},2)} &\equiv \left(\mathcal{W}_L^{-1} \otimes \mathcal{W}_L^{-1} \right) \circ \Delta_{L/\bar{L}}(\mathcal{W}(f)), \\
f_{(R/\bar{R},1)} \otimes f_{(R/\bar{R},2)} &\equiv \left(\mathcal{W}_R^{-1} \otimes \mathcal{W}_R^{-1} \right) \circ \Delta_{R/\bar{R}}(\mathcal{W}(f)).
\end{aligned} \tag{158}$$

Notice that in (156) and (157) the coordinate on top of the symbol for the action indicates the tensor factor the partial derivative acts upon.

There are several properties of the pairings in (153) and (154) worth recording here. As can be seen from their definition the values of the pairings are numbers. Due to this fact they should behave like scalars, i.e. the pairings

obey

$$
\begin{aligned}
h \triangleright \big\langle f(\partial^i), g(x^j) \big\rangle_{L,\bar{R}} &= \varepsilon(h) \big\langle f(\partial^i), g(x^j) \big\rangle_{L,\bar{R}}, \\
h \triangleright \big\langle f(\hat{\partial}^i), g(x^j) \big\rangle_{\bar{L},R} &= \varepsilon(h) \big\langle f(\hat{\partial}^i), g(x^j) \big\rangle_{\bar{L},R},
\end{aligned} \tag{159}
$$

$$
\begin{aligned}
h \triangleright \big\langle f(x^i), g(\partial^j) \big\rangle_{L,\bar{R}} &= \varepsilon(h) \big\langle f(x^i), g(\partial^j) \big\rangle_{L,\bar{R}}, \\
h \triangleright \big\langle f(x^i), g(\hat{\partial}^j) \big\rangle_{\bar{L},R} &= \varepsilon(h) \big\langle f(x^i), g(\hat{\partial}^j) \big\rangle_{\bar{L},R}.
\end{aligned} \tag{160}
$$

Furthermore, from the relations in (156) and (157) together with the definition of the dual pairings in (153) and (154) we find

$$
\begin{aligned}
\big\langle \partial^i \partial^j, f(x^k) \big\rangle_{L,\bar{R}} &= \big(\partial^i \partial^j \triangleright f(x^k) \big)\big|_{x^k=0} \\
&= \big(\partial^i \triangleright (\partial^j \triangleright f(x^k)) \big)\big|_{x^k=0} \\
&= \big\langle \partial^j, f_{(\bar{L},1)} \big\rangle_{L,\bar{R}} \big\langle \partial^i, f_{(\bar{L},2)} \big\rangle_{L,\bar{R}} \\
&= \big(\partial^j \overset{x}{\triangleright} [\partial^i \overset{y}{\triangleright} f(x^k \oplus_{\bar{L}} y^l)] \big)\big|_{x^k=y^l=0}, \\
\big\langle \hat{\partial}^i \hat{\partial}^j, f(x^k) \big\rangle_{\bar{L},R} &= \big\langle \hat{\partial}^j, f_{(L,1)} \big\rangle_{\bar{L},R} \big\langle \hat{\partial}^i, f_{(L,2)} \big\rangle_{\bar{L},R} \\
&= \big(\hat{\partial}^j \overset{x}{\triangleright} [\hat{\partial}^i \overset{y}{\triangleright} f(x^k \oplus_L y^l)] \big)\big|_{x^k=y^l=0},
\end{aligned} \tag{161}
$$

$$
\begin{aligned}
\big\langle f(x^k), \partial^j \partial^i \big\rangle_{L,\bar{R}} &= \big\langle f_{(R,1)}, \partial^i \big\rangle_{L,\bar{R}} \big\langle f_{(R,2)}, \partial^j \big\rangle_{L,\bar{R}} \\
&= \big([f(y^l \oplus_R x^k) \overset{y}{\triangleleft} \partial^j] \overset{x}{\triangleleft} \partial^i \big)\big|_{x^k=y^l=0}, \\
\big\langle f(x^k), \hat{\partial}^j \hat{\partial}^i \big\rangle_{\bar{L},R} &= \big\langle f_{(\bar{R},1)}, \hat{\partial}^i \big\rangle_{\bar{L},R} \big\langle f_{(\bar{R},2)}, \hat{\partial}^j \big\rangle_{\bar{L},R} \\
&= \big([f(y^l \oplus_{\bar{R}} x^k) \overset{y}{\triangleleft} \hat{\partial}^j] \overset{x}{\triangleleft} \hat{\partial}^i \big)\big|_{x^k=y^l=0}.
\end{aligned} \tag{162}
$$

In addition to this, we have

$$
\begin{aligned}
\big\langle 1, f(x^i) \big\rangle_{L,\bar{R}} &= f(x^i)|_{x^i=0}, & \big\langle 1, f(x^i) \big\rangle_{\bar{L},R} &= f(x^i)|_{x^i=0}, \\
\big\langle f(x^i), 1 \big\rangle_{L,\bar{R}} &= f(x^i)|_{x^i=0}, & \big\langle f(x^i), 1 \big\rangle_{\bar{L},R} &= f(x^i)|_{x^i=0},
\end{aligned} \tag{163}
$$

which again is a direct consequence of the definition of the dual pairings in (153) and (154).

There is another axiom the dual pairing has to fulfill. For this purpose

we perform the following calculation:

$$\begin{aligned}
&\big\langle f(\ominus_R\,\partial^i), g(x^j)\big\rangle_{L,\bar{R}} = \big(f(\ominus_R\,\partial^i)\triangleright g(x^j)\big)\big|_{x=0}\\
&= \big([\mathcal{R}^{-1}_{[1]}\triangleright g(x^j)]\triangleleft[(\mathcal{R}^{-1}_{[2]}\triangleright f(\partial^i)]\big)\big|_{x^j=0}\\
&= \big([\mathcal{R}^{-1}_{[1]}\triangleright g(x^j)]\,\bar{\triangleright}\,[\mathcal{R}^{-1}_{[2]}\triangleright f(\partial^i)]\big)\big|_{\partial^i=0}\\
&= \big(f(\partial^i)\,\bar{\triangleleft}\,g(\ominus_{\bar{L}}\,x^j)\big)\big|_{\partial^i=0} = \big\langle f(\partial^i), g(\ominus_{\bar{L}}\,x^j)\big\rangle_{L,\bar{R}},
\end{aligned}\tag{164}$$

where the second and fourth identity hold due to (146) and (147), respectively. With the same reasonings we can also show that

$$\begin{aligned}
\big\langle f(\ominus_{\bar{R}}\,\hat{\partial}^i), g(x^j)\big\rangle_{\bar{L},R} &= \big\langle f(\hat{\partial}^i), g(\ominus_L\,x^j)\big\rangle_{\bar{L},R},\\
\big\langle f(\partial^i), g(\ominus_{\bar{R}}\,x^j)\big\rangle_{L,\bar{R}} &= \big\langle f(\ominus_L\,\partial^i), g(x^j)\big\rangle_{L,\bar{R}},\\
\big\langle f(\hat{\partial}^i), g(\ominus_R\,x^j)\big\rangle_{\bar{L},R} &= \big\langle f(\ominus_{\bar{L}}\,\hat{\partial}^i), g(x^j)\big\rangle_{\bar{L},R}.
\end{aligned}\tag{165}$$

Interchanging the roles of coordinates and derivatives we additionally obtain

$$\begin{aligned}
\big\langle f(\ominus_R\,x^i), g(\partial^j)\big\rangle_{L,\bar{R}} &= \big\langle f(x^i), g(\ominus_{\bar{L}}\,\partial^j)\big\rangle_{L,\bar{R}},\\
\big\langle f(\ominus_{\bar{R}}\,x^i), g(\hat{\partial}^j)\big\rangle_{\bar{L},R} &= \big\langle f(x^i), g(\ominus_L\,\hat{\partial}^j)\big\rangle_{\bar{L},R},
\end{aligned}\tag{166}$$

$$\begin{aligned}
\big\langle f(x^i), g(\ominus_{\bar{R}}\,\partial^j)\big\rangle_{L,\bar{R}} &= \big\langle f(\ominus_L\,x^i), g(\partial^j)\big\rangle_{L,\bar{R}},\\
\big\langle f(x^i), g(\ominus_R\,\hat{\partial}^j)\big\rangle_{\bar{L},R} &= \big\langle f(\ominus_{\bar{L}}\,x^i), g(\hat{\partial}^j)\big\rangle_{\bar{L},R}.
\end{aligned}\tag{167}$$

4.2 Definition and basic properties of q-deformed exponentials

Let us make contact with another very important ingredient of braided tensor categories, i.e. an exponential. From an abstract point of view an exponential is nothing other than an object whose dualization is the pairing in (149) or (150). Thus, the exponential is given by the mapping

$$\exp : \mathbb{C} \longrightarrow \mathcal{A}_q \otimes \mathcal{A}_q^*, \quad \text{with} \quad \exp(\alpha) = \alpha \sum_a e_a \otimes f^a, \tag{168}$$

or

$$\exp' : \mathbb{C} \longrightarrow \mathcal{A}_q^* \otimes \mathcal{A}_q, \quad \text{with} \quad \exp'(\alpha) = \alpha \sum_a f^a \otimes e_a. \tag{169}$$

Again, the algebra isomorphism $\mathcal{W}$ enables us to introduce q-deformed exponentials that live on a tensor product of commutative algebras. More concretely, this is achieved by the expressions

$$\exp(x^i|\partial^j) \equiv \sum_a \mathcal{W}(e_a) \otimes \mathcal{W}(f^a), \tag{170}$$

and

$$\exp(\partial^i|x^j) \equiv \sum_a \mathcal{W}(f^a) \otimes \mathcal{W}(e_a). \tag{171}$$

If we want to derive explicit formulae for q-deformed exponentials it is our task to determine a basis of the coordinate algebra $\mathcal{A}_q$ being dual to a given one of the derivative algebra $\mathcal{A}_q^*$. Inserting the elements of the two bases into the formulae of (170) and (171) will then provide us with explicit expressions for q-deformed exponentials. It should be obvious that the two bases being dually paired depend on the choice of the pairing. Thus, each pairing in (153) and (154) leads to its own q-exponential:

$$\begin{aligned} \big\langle f(\partial^i), g(x^j) \big\rangle_{L,\bar{R}} \quad &\Rightarrow \quad \exp(x^i|\partial^j)_{\bar{R},L}, \\ \big\langle f(\hat{\partial}^i), g(x^j) \big\rangle_{\bar{L},R} \quad &\Rightarrow \quad \exp(x^i|\hat{\partial}^j)_{R,\bar{L}}, \end{aligned} \tag{172}$$

$$\begin{aligned} \big\langle f(x^i), g(\partial^j) \big\rangle_{L,\bar{R}} \quad &\Rightarrow \quad \exp(\partial^i|x^j)_{\bar{R},L}, \\ \big\langle f(x^i), g(\hat{\partial}^j) \big\rangle_{\bar{L},R} \quad &\Rightarrow \quad \exp(\hat{\partial}^i|x^j)_{R,\bar{L}}. \end{aligned} \tag{173}$$

In Ref. [40] we presented explicit formulae for q-deformed exponentials. Once again, the results for the two-dimensional quantum plane shall serve as an example. For the second exponential in (172) we found the expression

$$\exp(x^i|\hat{\partial}^j)_{R,\bar{L}} = \sum_{n_1,n_2=0}^{\infty} \frac{(x^1)^{n_1}(x^2)^{n_2} \otimes (\hat{\partial}_2)^{n_2}(\hat{\partial}_1)^{n_1}}{[[n_1]]_{q^{-2}}!\,[[n_2]]_{q^{-2}}!}. \tag{174}$$

Next, we wish to explain, in more detail, what it means that pairings and exponentials are dual to each other. To this aim, we should realize that the existence of two bases being dual to each other gives rise to completeness

relations of the form

$$u = \sum_a \langle u, e_a \rangle \, f^a \in \mathcal{A}_q^*, \quad y = \sum_a e_a \, \langle f^a, y \rangle \in \mathcal{A}_q, \tag{175}$$

and

$$u = \sum_a \langle e_a, u \rangle' \, f^a \in \mathcal{A}_q^*, \quad y = \sum_a e_a \, \langle y, f^a \rangle' \in \mathcal{A}_q. \tag{176}$$

From our definitions so far, it should be clear that the relations in (175) and (176) imply

$$\begin{aligned} \mathrm{id} &= \big(\langle .\,, . \rangle_{L,\bar{R}} \otimes \mathrm{id} \big) \circ \big(\mathrm{id} \otimes \exp(x^i | \partial^j)_{\bar{R},L} \big), \\ \mathrm{id} &= \big(\langle .\,, . \rangle_{\bar{L},R} \otimes \mathrm{id} \big) \circ \big(\mathrm{id} \otimes \exp(x^i | \hat{\partial}^j)_{R,\bar{L}} \big), \end{aligned} \tag{177}$$

$$\begin{aligned} \mathrm{id} &= \big(\mathrm{id} \otimes \langle .\,, . \rangle_{L,\bar{R}} \big) \circ \big(\exp(x^i | \partial^j)_{\bar{R},L} \otimes \mathrm{id} \big), \\ \mathrm{id} &= \big(\mathrm{id} \otimes \langle .\,, . \rangle_{\bar{L},R} \big) \circ \big(\exp(x^i | \hat{\partial}^j)_{R,\bar{L}} \otimes \mathrm{id} \big). \end{aligned} \tag{178}$$

Let us note that here and in what follows we limit ourselves to the exponentials of (172), since the relations for the other types of q-deformed exponentials are easily obtained by interchanging the roles for coordinates and derivatives.

Now, we would like to use the properties in (177) and (178) to derive the addition laws for q-exponentials. To this end, we show that the mapping in (170) has to be subject to

$$\begin{aligned} (\Delta \otimes \mathrm{id}) \exp &= \exp_{23} \exp_{13} = \sum_a e_a \otimes \Big(\sum_b e_b \otimes f^b \Big) f^a, \\ (\mathrm{id} \otimes \Delta) \exp &= \exp_{13} \exp_{12} = \sum_a e_a \Big(\sum_b e_b \otimes f^b \Big) \otimes f^a, \end{aligned} \tag{179}$$

where

$$\begin{aligned} \exp_{12}(\alpha) &\equiv \alpha \sum_a e_a \otimes f^a \otimes 1, \\ \exp_{23}(\alpha) &\equiv \alpha \sum_a 1 \otimes e_a \otimes f^a, \\ \exp_{13}(\alpha) &\equiv \alpha \sum_a e_a \otimes 1 \otimes f^a. \end{aligned} \tag{180}$$

It should be mentioned that the relations in (179) can be viewed as dualized versions of the identities [cf. (161) and (162)]

$$\begin{aligned}\big\langle uv, x\big\rangle &= \big\langle u, x_{(2)}\big\rangle\big\langle v, x_{(1)}\big\rangle,\\ \big\langle u, xy\big\rangle &= \big\langle u_{(2)}, x\big\rangle\big\langle u_{(1)}, y\big\rangle.\end{aligned} \tag{181}$$

For this reason the proof of the relations in (179) is mainly based on the identities in (181), as can be seen by the following calculations:

$$\begin{aligned}\exp_{23}\exp_{13} &= \sum_{a,b} e_a\otimes e_b\otimes f^b f^a\\ &\overset{(175)}{=} \sum_{a,b,c} e_a\otimes e_b\otimes \big\langle f^b f^a, e_c\big\rangle f^c\\ &\overset{(181)}{=} \sum_{a,b,c} e_a\otimes e_b\otimes \big\langle f^b, (e_c)_{(2)}\big\rangle\big\langle f^a, (e_c)_{(1)}\big\rangle f^c\\ &\overset{(175)}{=} \sum_{c} (e_c)_{(1)}\otimes (e_c)_{(2)}\otimes f^c = (\Delta\otimes\mathrm{id})\exp,\end{aligned} \tag{182}$$

and

$$\begin{aligned}\exp_{13}\exp_{12} &= \sum_{a,b} e_a e_b\otimes f^b\otimes f^a\\ &\overset{(175)}{=} \sum_{a,b,c} e_c\big\langle e_a e_b, f^c\big\rangle'\otimes f^b\otimes f^a\\ &\overset{(181)}{=} \sum_{a,b,c} e_c\big\langle e_a, f^c_{(2)}\big\rangle'\big\langle e_b, f^c_{(1)}\big\rangle'\otimes f^b\otimes f^a\\ &\overset{(175)}{=} \sum_{c} e_c\otimes f^c_{(1)}\otimes f^c_{(2)} = (\mathrm{id}\otimes\Delta)\exp.\end{aligned} \tag{183}$$

It should be obvious that the relations in (179) carry over to q-deformed exponentials on commutative algebras. This way, they can be reformulated as

$$\exp(x^i\oplus_L y^j|\hat\partial^k)_{R,\bar L} = \exp(x^i|\exp(y^j|\hat\partial^k)_{R,\bar L}\overset{\partial}{\circledast}\hat\partial^l)_{R,\bar L},$$

$$\exp(x^i \oplus_{\bar{L}} y^j|\partial^k)_{\bar{R},L} = \exp(x^i|\exp(y^j|\partial^k)_{\bar{R},L} \overset{\partial}{\circledast} \partial^l)_{\bar{R},L}, \tag{184}$$

$$\begin{aligned}\exp(x^k|\hat{\partial}^j \oplus_{\bar{R}} \hat{\partial}'^i)_{R,\bar{L}} &= \exp(x^l \overset{x}{\circledast} \exp(x^k|\hat{\partial}^j)_{R,\bar{L}}|\hat{\partial}'^i)_{R,\bar{L}},\\ \exp(x^k|\partial^j \oplus_{R} \partial'^i)_{\bar{R},L} &= \exp(x^l \overset{x}{\circledast} \exp(x^k|\partial^j)_{\bar{R},L}|\partial'^i)_{\bar{R},L}.\end{aligned} \tag{185}$$

To understand the physical meaning of q-exponentials it is recorded that they can be regarded as q-analogs of classical plane waves. In other words, q-exponentials are in some sense eigenfunctions of partial derivatives, since they obey

$$\begin{aligned}\partial^i \overset{x}{\triangleright} \exp(x^k|\partial^l)_{\bar{R},L} &= \exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial}{\circledast} \partial^i,\\ \hat{\partial}^i \overset{x}{\triangleright} \exp(x^k|\hat{\partial}^l)_{R,\bar{L}} &= \exp(x^k|\hat{\partial}^l)_{R,\bar{L}} \overset{\partial}{\circledast} \hat{\partial}^i,\end{aligned} \tag{186}$$

$$\begin{aligned}\exp(\partial^l|x^k)_{\bar{R},L} \overset{x}{\triangleleft} \partial^i &= \partial^i \overset{\partial}{\circledast} \exp(\partial^l|x^k)_{\bar{R},L},\\ \exp(\hat{\partial}^l|x^k)_{R,\bar{L}} \overset{x}{\triangleleft} \hat{\partial}^i &= \hat{\partial}^i \overset{\partial}{\circledast} \exp(\hat{\partial}^l|x^k)_{R,\bar{L}},\end{aligned} \tag{187}$$

and

$$\begin{aligned}\exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial}{\triangleleft} x^i &= x^i \overset{x}{\circledast} \exp(x^k|\partial^l)_{\bar{R},L},\\ \exp(x^k|\hat{\partial}^l)_{R,\bar{L}} \overset{\partial}{\triangleleft} x^i &= x^i \overset{x}{\circledast} \exp(x^k|\hat{\partial}^l)_{R,\bar{L}},\end{aligned} \tag{188}$$

$$\begin{aligned}x^i \overset{\partial}{\triangleright} \exp(\partial^l|x^k)_{\bar{R},L} &= \exp(\partial^l|x^k)_{\bar{R},L} \overset{x}{\circledast} x^i,\\ x^i \overset{\partial}{\triangleright} \exp(\hat{\partial}^l|x^k)_{R,\bar{L}} &= \exp(\hat{\partial}^l|x^k)_{R,\bar{L}} \overset{x}{\circledast} x^i.\end{aligned} \tag{189}$$

For a proof of the above identities we need the coregular actions in (156) and (157), which more abstractly take the form

$$\begin{aligned}\partial^i \triangleright f &= \left\langle \partial^i, f_{(1)} \right\rangle f_{(2)},\\ f \triangleleft \partial^i &= f_{(2)} \left\langle f_{(1)}, \partial^i \right\rangle.\end{aligned} \tag{190}$$

With these identities at hand a straightforward computation yields

$$\partial^i \triangleright \exp(x^k|\partial^l) = \sum_a \left(\partial^i \triangleright \mathcal{W}(e_a)\right) \otimes \mathcal{W}(f^a)$$

$$\overset{(190)}{=} \sum_a \big\langle \partial^i, \mathcal{W}(e_a)_{(1)} \big\rangle \mathcal{W}(e_a)_{(2)} \otimes \mathcal{W}(f^a)$$
$$\overset{(179)}{=} \sum_{a,b} \big\langle \partial^i, \mathcal{W}(e_a) \big\rangle \mathcal{W}(e_b) \otimes \mathcal{W}(f^b)\mathcal{W}(f^a)$$
$$\overset{(175)}{=} \sum_a \mathcal{W}(e_a) \otimes \mathcal{W}(f^a)\partial^i = \exp(x^k|\partial^l) \overset{\partial}{\circledast} \partial^i. \tag{191}$$

Through a slight modification to these arguments one can prove the corresponding identities concerning right actions of x^i.

For the sake of completeness let us note that our q-exponentials are normalized in such a way that

$$\exp(x^i|\partial^j)_{\bar{R},L}|_{x^i=0} = \exp(x^i|\partial^j)_{\bar{R},L}|_{\partial^j=0} = 1,$$
$$\exp(x^i|\hat{\partial}^j)_{R,\bar{L}}|_{x^i=0} = \exp(x^i|\hat{\partial}^j)_{R,\bar{L}}|_{\partial^j=0} = 1. \tag{192}$$

These equalities result from

$$\exp(x^i|\partial^j)|_{x^i=0} \overset{(163)}{=} \sum_a \big\langle 1, \mathcal{W}(e_a) \big\rangle \otimes \mathcal{W}(f^a)$$
$$\overset{(175)}{=} 1 \otimes 1 = 1, \tag{193}$$

and

$$\exp(x^i|\partial^j)|_{x^i=0} \overset{(163)}{=} \sum_a \mathcal{W}(e_a) \otimes \big\langle \mathcal{W}(f^a), 1 \big\rangle$$
$$\overset{(175)}{=} 1 \otimes 1 = 1. \tag{194}$$

It is well-known that the generators of the quantum algebras describing the underlying symmetry can be realized within the Weyl-algebras $\mathcal{A}_q^* \otimes \mathcal{A}_q$ and $\mathcal{A}_q \otimes \mathcal{A}_q^*$, i.e. they can be represented in terms of coordinates and partial derivatives [18, 71]. Due to this fact, it immediately follows from (186) and (188) that q-deformed exponentials satisfy

$$h \overset{x}{\triangleright} \exp(x^i|\partial^j)_{\bar{R},L} = \exp(x^i|\partial^j)_{\bar{R},L} \overset{\partial}{\triangleleft} h,$$
$$h \overset{x}{\triangleright} \exp(x^i|\hat{\partial}^j)_{R,\bar{L}} = \exp(x^i|\hat{\partial}^j)_{R,\bar{L}} \overset{\partial}{\triangleleft} h, \tag{195}$$

where h stands for an element of the symmetry algebra. With these relations at hand we can now prove that q-deformed exponentials behave like scalars:

$$\begin{aligned}
& h \triangleright \big(\exp(x^i|\partial^j) \big) \\
&= \sum_a \big(h_{(1)} \triangleright \mathcal{W}(e_a) \big) \otimes \big(h_{(2)} \triangleright \mathcal{W}(f^a) \big) \\
&\overset{(195)}{=} \sum_a \mathcal{W}(e_a) \otimes \big(h_{(2)} \triangleright (\mathcal{W}(f^a) \triangleleft h_{(1)}) \big) \\
&= \sum_a \mathcal{W}(e_a) \otimes \big(h_{(2)} S^{-1}(h_{(1)}) \triangleright \mathcal{W}(f^a) \big) \\
&= \varepsilon(h) \sum_a \mathcal{W}(e_a) \otimes \mathcal{W}(f^a) = \varepsilon(h) \exp(x^i|\partial^j),
\end{aligned} \tag{196}$$

$$\begin{aligned}
& \big(\exp(x^i|\partial^j) \big) \triangleleft h \\
&= \sum_a \big(\mathcal{W}(e_a) \triangleleft h_{(2)} \big) \otimes \big(\mathcal{W}(f^a) \triangleleft h_{(1)} \big) \\
&\overset{(195)}{=} \sum_a \big((h_{(1)} \triangleright \mathcal{W}(e_a)) \triangleleft h_{(2)} \big) \otimes \mathcal{W}(f^a) \\
&= \sum_a \big(\mathcal{W}(e_a) \triangleleft S(h_{(1)}) h_{(2)} \big) \otimes \mathcal{W}(f^a) \\
&= \varepsilon(h) \sum_a \mathcal{W}(e_a) \otimes \mathcal{W}(f^a) = \varepsilon(h) \exp(x^i|\partial^j).
\end{aligned} \tag{197}$$

Next, we come to the dualized versions of the relations in (164)-(167). They are given by

$$\begin{aligned}
\exp(\ominus_{\bar{L}}\, x^i|\partial^j)_{\bar{R},L} &= \exp(x^i|\ominus_R\, \partial^j)_{\bar{R},L}, \\
\exp(\ominus_L\, x^i|\partial^j)_{R,\bar{L}} &= \exp(x^i|\ominus_{\bar{R}}\, \partial^j)_{R,\bar{L}},
\end{aligned} \tag{198}$$

$$\begin{aligned}
\exp(\ominus_{\bar{R}}\, x^i|\partial^j)_{\bar{R},L} &= \exp(x^i|\ominus_L\, \partial^j)_{\bar{R},L}, \\
\exp(\ominus_R\, x^i|\partial^j)_{R,\bar{L}} &= \exp(x^i|\ominus_{\bar{L}}\, \partial^j)_{R,\bar{L}}.
\end{aligned} \tag{199}$$

To prove them we first perform calculations like the following one:

$$\sum_a \big\langle g(\partial^i), \bar{S}(e_a) \big\rangle f^a$$

$$
\begin{aligned}
&= \sum_a \big\langle S^{-1}(g(\partial^i)), e_a \big\rangle f^a \\
&= S^{-1}(g(\partial^i)) = \sum_a \big\langle g(\partial^i), e_a \big\rangle S^{-1}(f^a),
\end{aligned} \tag{200}
$$

where $\bar{S}$ and S^{-1} respectively denote conjugate and inverse of the antipode of the braided Hopf algebra $\mathcal{A}$ [cf. (55) and (56)]. For the first identity in (200) we applied the result of (164), while for the second and third equality we made use of (175). Realizing that the pairing is non-degenerate we can finally read off from (200) that

$$
\sum_a \bar{S}(e_a) \otimes f^a = \sum_a e_a \otimes S^{-1}(f^a), \tag{201}
$$

which is equivalent to the first relation in (198). Similar considerations lead to the other relations in (198) and (199).

It should also be mentioned that in some sense the exponentials in (198) are inverse to those introduced in (172), since they obey the identities

$$
\begin{aligned}
&\exp(x^i \overset{x}{\circledast} \exp(\ominus_{\bar{L}}\, x^j|\partial^k)_{\bar{R},L} \overset{\partial}{\circledast} \partial^l)_{\bar{R},L} \\
&\overset{(184)}{=} \exp(x^i \oplus_{\bar{L}} (\ominus_{\bar{L}}\, x^j)|\partial^k)_{\bar{R},L} \\
&\overset{(65)}{=} \exp(x^i|\partial^k)_{\bar{R},L}|_{x^i=0} = 1 \\
&\overset{(65)}{=} \exp((\ominus_{\bar{L}}\, x^j) \oplus_{\bar{L}} x^i|\partial^k)_{\bar{R},L} \\
&\overset{(184)}{=} \exp((\ominus_{\bar{L}}\, x^j) \overset{x}{\circledast} \exp(x^i|\partial^k)_{\bar{R},L} \overset{\partial}{\circledast} \partial^l)_{\bar{R},L},
\end{aligned} \tag{202}
$$

and

$$
\begin{aligned}
&\exp(x^i \overset{x}{\circledast} \exp(x^j|\ominus_R\, \partial^k)_{\bar{R},L} \overset{\partial}{\circledast} \partial^l)_{\bar{R},L} \\
&\overset{(185)}{=} \exp(x^i|(\ominus_R\, \partial^k) \oplus_R \partial^l)_{\bar{R},L} \\
&\overset{(66)}{=} \exp(x^i|\partial^k)_{\bar{R},L}|_{\partial^k=0} = 1 \\
&\overset{(66)}{=} \exp(x^i|\partial^k \oplus_R (\ominus_R\, \partial^l))_{\bar{R},L} \\
&\overset{(185)}{=} \exp(x^i \overset{x}{\circledast} \exp(x^j|\partial^k)_{\bar{R},L} \overset{\partial}{\circledast} (\ominus_R\, \partial^l))_{\bar{R},L}.
\end{aligned} \tag{203}
$$

Before proceeding further we would like to examine wether the inverse exponentials also show the property of being eigenfunctions of partial derivatives. This is indeed the case, but compared to (186) in a somewhat different way, as can be shown by the following calculation:

$$\begin{aligned}
&\sum_a (\partial^i \triangleright e_a) \otimes S^{-1}(f^a) \\
&= \sum_a e_a \otimes S^{-1}(f^a \partial^i) \\
&= -\sum_a e_a \otimes \partial^j S^{-1}(f^a \triangleleft \mathcal{L}^i_j) \\
&= -\sum_a (1 \otimes \partial^k)\big((e_a \triangleleft \mathcal{L}^j_k) \otimes S^{-1}(f^a \triangleleft \mathcal{L}^i_j)\big) \\
&= -(1 \otimes \partial^i) \sum_a e_a \otimes S^{-1}(f^a) \\
&= -\sum_a (\mathcal{L}^i_j \triangleright e_a) \otimes \partial^j S^{-1}(f^a). \qquad (204)
\end{aligned}$$

The first equality is nothing else than (186) and the second equality uses (50) together with $S^{-1}(\partial^i) = -\partial^i$. For the third equality we commute the partial derivative to the left of the exponential. The fourth equality results from the fact that exponentials behave like scalars [cf. (196) and (197)]. Thus, exponentials are invariant under the action of $\mathcal{L}^i_j$ (notice that for the entries of the L-matrix we have $\Delta(\mathcal{L}^i_j) = \mathcal{L}^i_k \otimes \mathcal{L}^k_j$ and $\varepsilon(\mathcal{L}^i_j) = \delta^i_j$). It should be clear that on commutative algebras the result in (204) reads as

$$\partial^i \overset{x}{\triangleright} \exp(x^k|\ominus_R \partial^l)_{\bar{R},L} = -\partial^i \overset{\partial|x}{\odot}_L \exp(x^k|\ominus_R \partial^l)_{\bar{R},L}, \qquad (205)$$

where the factors of the braided product are indicated by the variables on top of the corresponding symbol. Likewise, we have

$$\hat{\partial}^i \overset{x}{\bar{\triangleright}} \exp(x^k|\ominus_{\bar{R}} \hat{\partial}^l)_{R,\bar{L}} = -\hat{\partial}^i \overset{\partial|x}{\odot}_{\bar{L}} \exp(x^k|\ominus_{\bar{R}} \hat{\partial}^l)_{R,\bar{L}}, \qquad (206)$$

and

$$\exp(x^k|\ominus_R \partial^l)_{\bar{R},L} \overset{\partial}{\triangleleft} x^i = -\exp(x^k|\ominus_R \partial^l)_{\bar{R},L} \overset{\partial|x}{\odot}_{\bar{R}} x^i,$$

$$\exp(x^k|\ominus_{\bar{R}}\,\hat{\partial}^l)_{R,\bar{L}}\overset{\partial}{\triangleleft}x^i=-\exp(x^k|\ominus_{\bar{R}}\,\hat{\partial}^l)_{R,\bar{L}}\overset{\partial|x}{\odot}{}_R\,x^i. \tag{207}$$

So far it seems that the inverse exponentials give another set of plane-waves, since the identities in (205)-(207) differ from those in (186) and (188). Remarkably, the inverse exponentials can be identified with the exponentials introduced in (172). For this to become clear, we show that the inverse exponentials satisfy the same relations as the original ones. With the help of the identities in (141), we get

$$\begin{aligned}&\sum_a(\partial^i\triangleright\bar{S}^{-1}(e_a))\otimes f^a\\&=-\sum_a S^{-1}(\hat{\partial}^i\,\bar{\triangleright}\,e_a)\otimes f^a\\&=-\sum_a\bar{S}^{-1}(e_a)\otimes(f^a\hat{\partial}^i)\end{aligned} \tag{208}$$

being tantamount to

$$\partial^i\overset{x}{\triangleright}\exp(\ominus_{\bar{R}}\,x^k|\hat{\partial}^l)_{R,\bar{L}}=\exp(\ominus_{\bar{R}}\,x^k|\hat{\partial}^l)_{R,\bar{L}}\overset{\partial}{\circledast}(-\hat{\partial}^i). \tag{209}$$

The point now is that the exponentials in (172) are completely characterized by their relations in (186). Thus, comparing the result of (208) to the relations in (186) yields

$$\exp(\ominus_{\bar{R}}\,x^i|\hat{\partial}^j)_{R,\bar{L}}=\exp(x^i|-\hat{\partial}^j)_{\bar{R},L}\equiv\exp(x^i|\partial^j)_{\bar{R},L}|_{\partial^j\to-\hat{\partial}^j}, \tag{210}$$

which by virtue of $f(\ominus_{\bar{L}}(\ominus_{\bar{R}}\,x^i))=f(x^i)$ leads to the equality

$$\exp(\ominus_{\bar{L}}x^i|\partial^j)_{\bar{R},L}=\exp(x^i|-\partial^j)_{R,\bar{L}}\equiv\exp(x^i|\hat{\partial}^j)_{R,\bar{L}}|_{\hat{\partial}^j\to-\partial^j}. \tag{211}$$

In the same way we can verify that

$$\exp(\ominus_R x^i|\partial^j)_{\bar{R},L}=\exp(x^i|-\partial^j)_{R,\bar{L}}\equiv\exp(x^i|\hat{\partial}^j)_{R,\bar{L}}|_{\hat{\partial}^j\to-\partial^j}, \tag{212}$$

and

$$\exp(\ominus_L x^i|\hat{\partial}^j)_{R,\bar{L}}=\exp(x^i|-\hat{\partial}^j)_{\bar{R},L}\equiv\exp(x^i|\partial^j)_{\bar{R},L}|_{\partial^j\to-\hat{\partial}^j}. \tag{213}$$

Notice that for the Manin plane the minus sign in the expressions (209)-(213)

must be dropped.

4.3 q-Deformed analogs of Taylor rules

Now, we come to a very useful application of our q-deformed exponentials, i.e. q-deformed Taylor rules. First of all, we should note that our exponentials generate q-deformed translations in the following sense:

$$\begin{aligned}\exp(x^i|\partial^j)_{\bar{R},L} \overset{\partial|y}{\triangleright} g(y^k) &= g(x^i \oplus_{\bar{L}} y^k),\\ \exp(x^i|\hat{\partial}^j)_{R,\bar{L}} \overset{\partial|y}{\triangleright} g(y^k) &= g(x^i \oplus_{L} y^k),\end{aligned} \tag{214}$$

and

$$\begin{aligned} g(y^k) \overset{y|\partial}{\triangleleft} \exp(\partial^j|x^i)_{\bar{R},L} &= g(y^k \oplus_{R} x^i),\\ g(y^k) \overset{y|\partial}{\triangleleft} \exp(\hat{\partial}^j|x^i)_{R,\bar{L}} &= g(y^k \oplus_{\bar{R}} x^i),\end{aligned} \tag{215}$$

where we have extended our notation in order to indicate the acting object. The above identities can be proved by straightforward calculations, i.e.

$$\begin{aligned}\exp(x^i|\partial^j)_{\bar{R},L} \overset{\partial|y}{\triangleright} g(y^k) &= \sum_a \mathcal{W}(e_a) \otimes \big\langle \mathcal{W}(f^a), g(y^k)_{(\bar{L},1)} \big\rangle_{L,\bar{R}}\, g(y^k)_{(\bar{L},2)}\\ &= \sum_a \mathcal{W}(e_a) \otimes \varepsilon\big(\mathcal{W}(f^a) \,\bar{\triangleleft}\, g(y^k)_{(\bar{L},1)}\big) g(y^k)_{(\bar{L},2)}\\ &= \sum_a \big(g(x^i)_{(\bar{L},1)} \overset{x}{\circledast} \mathcal{W}(e_a)\big) \otimes \varepsilon\left(\mathcal{W}(f^a)\right) g(y^k)_{(\bar{L},2)}\\ &= g(x^i)_{(\bar{L},1)} \otimes g(y^k)_{(\bar{L},2)} = g(x^i \oplus_{\bar{L}} y^k),\end{aligned} \tag{216}$$

and

$$\begin{aligned} g(y^k) \overset{y|\partial}{\triangleleft} \exp(\partial^j|x^i)_{\bar{R},L} &= \sum_a g(y^k)_{(R,1)} \big\langle g(y^k)_{(R,2)}, \mathcal{W}(f^a) \big\rangle_{L,\bar{R}} \otimes \mathcal{W}(e_a)\\ &= \sum_a g(y^k)_{(R,1)}\, \varepsilon\big(g(y^k)_{(R,2)} \triangleright \mathcal{W}(f^a)\big) \otimes \mathcal{W}(e_a)\\ &= \sum_a g(y^k)_{(R,1)}\, \varepsilon\left(\mathcal{W}(f^a)\right) \otimes \big(\mathcal{W}(e_a) \circledast g(x^i)_{(R,2)}\big)\end{aligned}$$

$$= g(y^k)_{(R,1)} \otimes g(x^i)_{(R,2)} = g(y^k \oplus_R x^i). \tag{217}$$

For the first step we have rewritten the action of the partial derivatives by (156) and (157). The second equality is the definition of the pairing. The third and fourth equality respectively use the properties in (186) and the normalization conditions in (192).

It is very instructive to expand the q-exponentials in (214) and (215) up to terms linear in x^i. In doing so, we obtain

$$\begin{aligned} f(x^i \oplus_L y^j) &= 1 \otimes f(y^j) + x^k \otimes \hat{\partial}_k \,\bar{\triangleright}\, f(y^j) + O(x^2), \\ f(x^i \oplus_{\bar{L}} y^j) &= 1 \otimes f(y^j) + x^k \otimes \partial_k \triangleright f(y^j) + O(x^2), \end{aligned} \tag{218}$$

$$\begin{aligned} f(y^i \oplus_{\bar{R}} x^j) &= f(y^i) \otimes 1 - f(y^i) \triangleleft \hat{\partial}^k \otimes x_k + O(x^2), \\ f(y^i \oplus_R x^j) &= f(y^i) \otimes 1 - f(y^i) \,\bar{\triangleleft}\, \partial^k \otimes x_k + O(x^2). \end{aligned} \tag{219}$$

As one expects we see that the partial derivatives indeed generate infinitesimal translations. Furthermore, the above relations give a hint to another possibility for defining the action of partial derivatives. In analogy to the undeformed situation, we have

$$\begin{aligned} \partial_i \triangleright f(x^j) &= \lim_{y^k \to 0} \frac{f(y^k \oplus_{\bar{L}} x^j) - f(x^j)}{y^i}, \\ \hat{\partial}_i \,\bar{\triangleright}\, f(x^j) &= \lim_{y^k \to 0} \frac{f(y^k \oplus_L x^j) - f(x^j)}{y^i}, \end{aligned} \tag{220}$$

$$\begin{aligned} f(x^j) \,\bar{\triangleleft}\, \partial^i &= \lim_{y^k \to 0} \frac{f(x^j) - f(x^j \oplus_R y^k)}{y_i}, \\ f(x^j) \triangleleft \hat{\partial}^i &= \lim_{y^k \to 0} \frac{f(x^j) - f(x^j \oplus_{\bar{R}} y^k)}{y_i}. \end{aligned} \tag{221}$$

Now, we are in a position to write down q-deformed Taylor rules, which allow us to determine the value of a function at a certain point from its values at a different point. From what we have done so far, it is not very difficult to convince oneself that we have

$$\begin{aligned} f(y^i) &= f((y^i \oplus_{\bar{L}} (\ominus_{\bar{L}} x^j)) \oplus_{\bar{L}} x^k) \\ &= \exp(y^i \oplus_{\bar{L}} (\ominus_{\bar{L}} x^j) | \partial^l)_{\bar{R},L} \overset{\partial|x}{\triangleright} f(x^k), \end{aligned}$$

$$
\begin{aligned}
f(y^i) &= f((y^i \oplus_L (\ominus_L x^j)) \oplus_L x^k) \\
&= \exp(y^i \oplus_L (\ominus_L x^j)|\hat{\partial}^l)_{R,\bar{L}} \overset{\partial|x}{\triangleright} f(x^k),
\end{aligned} \tag{222}
$$

and

$$
\begin{aligned}
f(y^i) &= f(x^k \oplus_R ((\ominus_R x^j) \oplus_R y^i)) \\
&= f(x^k) \overset{x|\partial}{\triangleleft} \exp(\partial^l|(\ominus_R x^j) \oplus_R y^i)_{\bar{R},L}, \\
f(y^i) &= f(x^k \oplus_{\bar{R}} ((\ominus_{\bar{R}} x^j) \oplus_{\bar{R}} y^i)) \\
&= f(x^k) \overset{x|\partial}{\bar{\triangleleft}} \exp(\hat{\partial}^l|(\ominus_{\bar{R}} x^j) \oplus_{\bar{R}} y^i)_{R,\bar{L}}.
\end{aligned} \tag{223}
$$

4.4 Conjugation properties and crossing symmetries

Next, we turn to the conjugation properties of dual pairings and q-deformed exponentials. We start our considerations with the following calculation:

$$
\begin{aligned}
\overline{\big\langle f(\partial^i), g(x^j)\big\rangle}_{L,\bar{R}} &= \overline{\varepsilon(f(\partial^i) \triangleright g(x^j))} = \varepsilon(\overline{f(\partial^i) \triangleright g(x^j)}) \\
&= \varepsilon(\overline{g(x^i)} \,\bar{\triangleleft}\, \overline{f(\partial^j)}) = \big\langle \overline{g(x^i)}, \overline{f(\partial^j)}\big\rangle_{L,\bar{R}}.
\end{aligned} \tag{224}
$$

In the same way, we can verify

$$
\overline{\big\langle f(\hat{\partial}^i), g(x^j)\big\rangle}_{\bar{L},R} = \big\langle \overline{g(x^j)}, \overline{f(\hat{\partial}^i)}\big\rangle_{\bar{L},R}, \tag{225}
$$

and

$$
\begin{aligned}
\overline{\big\langle f(x^i), g(\partial^j)\big\rangle}_{L,\bar{R}} &= \big\langle \overline{g(\partial^j)}, \overline{f(x^i)}\big\rangle_{L,\bar{R}}, \\
\overline{\big\langle f(x^i), g(\hat{\partial}^j)\big\rangle}_{\bar{L},R} &= \big\langle \overline{g(\hat{\partial}^j)}, \overline{f(x^i)}\big\rangle_{\bar{L},R}.
\end{aligned} \tag{226}
$$

In order to derive the conjugation properties for q-deformed exponentials, let us recall that q-deformed exponentials are completely characterized by the equations in (186) and (188). Thus, the conjugation properties of q-deformed exponentials can easily be read off if we compare these equations with their conjugate versions. For this to become more clear we proceed as follows:

$$
\overline{\partial^k \overset{x}{\triangleright} \exp(x^i|\partial^j)_{\bar{R},L}} = \overline{\exp(x^i|\partial^j)_{\bar{R},L} \overset{\partial}{\circledast} \partial^k},
$$

$$\Rightarrow \quad \overline{\exp(x^i|\partial^j)}_{\bar{R},L} \overset{x}{\triangleleft} \partial^k = \partial^k \overset{\partial}{\circledast} \overline{\exp(x^i|\partial^j)}_{\bar{R},L}. \tag{227}$$

Comparing the last result with

$$\exp(\partial^j|x^i)_{\bar{R},L} \overset{x}{\triangleleft} \partial^k = \partial^k \overset{\partial}{\circledast} \exp(\partial^j|x^i)_{\bar{R},L} \tag{228}$$

yields the identification

$$\overline{\exp(x^i|\partial^j)}_{\bar{R},L} = \exp(\partial^j|x^i)_{\bar{R},L}. \tag{229}$$

Similar considerations additionally give

$$\overline{\exp(x^i|\hat{\partial}^j)}_{R,\bar{L}} = \exp(\hat{\partial}^j|x^i)_{R,\bar{L}}, \tag{230}$$

and

$$\begin{aligned} \overline{\exp(\partial^i|x^j)}_{\bar{R},L} &= \exp(x^j|\partial^i)_{\bar{R},L}, \\ \overline{\exp(\hat{\partial}^i|x^j)}_{R,\bar{L}} &= \exp(x^j|\hat{\partial}^i)_{R,\bar{L}}. \end{aligned} \tag{231}$$

Now we come to the crossing symmetries for pairings and exponentials. As we know there are different versions of dual pairings and q-deformed exponentials. Again, the crossing symmetries are responsible for the fact that the expressions for the different versions of dual pairings and q-deformed exponentials can be transformed into each other by applying some simple substitutions. Fig. 3 shows the concrete relationship between the different types of pairings and exponentials. Again, we illustrate the transition rules by an example referring to the Manin plane [40]:

$$\begin{aligned} & \big\langle (\hat{\partial}_2)^{n_2} (\hat{\partial}_1)^{n_1}, (x^1)^{m_1} (x^2)^{m_2} \big\rangle_{\bar{L},R} \\ & = \delta_{m_1,n_1} \delta_{m_2,n_2} \, [[n_1]]_{q^{-2}}! \, [[n_2]]_{q^{-2}}! \\ \longleftrightarrow \quad & \big\langle (\partial_1)^{n_1} (\partial_2)^{n_2}, (x^2)^{m_2} (x^1)^{m_1} \big\rangle_{L,\bar{R}} \\ & = \delta_{m_1,n_1} \delta_{m_2,n_2} \, [[n_1]]_{q^2}! \, [[n_2]]_{q^2}! \\ \longleftrightarrow \quad & \big\langle (x_1)^{n_1} (x_2)^{n_2}, (-\partial^2)^{m_2} (-\partial^1)^{m_1} \big\rangle_{L,\bar{R}} \\ & = \delta_{m_1,n_1} \delta_{m_2,n_2} \, [[n_1]]_{q^2}! \, [[n_2]]_{q^2}!, \end{aligned} \tag{232}$$




Figure 3: Crossing-symmetries for braided products and q-translations

and

$$\exp(x^i|\hat{\partial}^j)_{R,\bar{L}} = \sum_{n_1,n_2=0}^{\infty} \frac{(x^1)^{n_1}(x^2)^{n_2} \otimes (\hat{\partial}_2)^{n_2}(\hat{\partial}_1)^{n_1}}{[[n_1]]_{q^{-2}}![[n_2]]_{q^{-2}}!}$$

$$\longleftrightarrow \quad \exp(x^i|\partial^j)_{\bar{R},L} = \sum_{n_1,n_2=0}^{\infty} \frac{(x^2)^{n_2}(x^1)^{n_1} \otimes (\partial_1)^{n_1}(\partial_2)^{n_2}}{[[n_1]]_{q^2}![[n_2]]_{q^2}!}$$

$$\longleftrightarrow \quad \exp(\partial^i|x^j)_{\bar{R},L} = \sum_{n_1,n_2=0}^{\infty} \frac{(-\partial^2)^{n_2}(-\partial^1)^{n_1} \otimes (x_1)^{n_1}(x_2)^{n_2}}{[[n_1]]_{q^2}![[n_2]]_{q^2}!}. \qquad (233)$$

We would like to close this section with a comment on q-deformed exponentials in terms of momentum variables. It should be clear that there is no reason to prevent us from substituting $\mathrm{i}^{-1}p^k$ for ∂^k in the expressions for q-deformed exponentials.

5 Integration on quantum spaces

In this section we introduce a powerful concept of integration on q-deformed quantum spaces. We start with presenting the basic ideas and illustrate them by the results for the two-dimensional quantum plane. We shall then explore the main features of q-deformed integrals. For other concepts of integration on quantum spaces we refer the reader to Refs. [66–69].

5.1 Inverse partial derivatives and their representations

Let us recall that integrals can be seen as operations being inverse to partial derivatives. Thus, we first try to extend the algebra of partial derivatives by introducing inverse elements. In the case of the two-dimensional quantum plane the algebra of partial derivatives is now characterized by the following relations:

$$\partial^2\partial^1 = q^{-1}\partial^1\partial^2, \tag{234}$$

$$\begin{aligned} \partial^i(\partial^i)^{-1} &= (\partial^i)^{-1}\partial^i = 1, \quad i = 1,2, \\ \partial^2(\partial^1)^{-1} &= q(\partial^1)^{-1}\partial^2, \\ \partial^1(\partial^2)^{-1} &= q(\partial^2)^{-1}\partial^1, \end{aligned} \tag{235}$$

$$(\partial^2)^{-1}(\partial^1)^{-1} = q^{-1}(\partial^1)^{-1}(\partial^2)^{-1}. \tag{236}$$

It should be mentioned that the commutation relations involving inverse partial derivatives can be derived most easily by means of the operator expressions for star products. To understand this more properly we give the following calculation holding for the Manin plane:

$$\begin{aligned} (X^i)^{\pm 1}(X^j)^{\pm 1} &= \mathcal{W}\big(\mathcal{W}^{-1}((X^i)^{\pm 1}) \circledast \mathcal{W}^{-1}((X^j)^{\pm 1})\big) \\ &= \mathcal{W}\big((x^i)^{\pm 1} \circledast (x^j)^{\pm 1}\big) \\ &= \mathcal{W}\big(\big[q^{-\hat{n}_{x^2}\hat{n}_{y^1}}(x^i)^{\pm 1}(y^j)^{\pm 1}\big]_{y^j \to x^i}\big) \\ &= q^{-(\pm\delta_{i2})(\pm\delta_{j1})}\mathcal{W}\big((x^i)^{\pm 1}(x^j)^{\pm 1}\big). \end{aligned} \tag{237}$$

For the sake of completeness we would like to mention that inverse partial derivatives show well-defined transformation properties. In the case of the

two-dimensional quantum plane, for example, the relations (234)-(236) together with the commutation relations between $U_q(su_2)$-generators and partial derivatives imply

$$\begin{aligned} T^+(\partial^1)^{-1} &= q^{-1}(\partial^1)^{-1}T^+ - q^{-1}(\partial^1)^{-2}\partial^2, \\ T^+(\partial^2)^{-1} &= q(\partial^2)^{-1}T^+, \end{aligned} \tag{238}$$

$$\begin{aligned} T^-(\partial^1)^{-1} &= q^{-1}(\partial^1)^{-1}T^-, \\ T^-(\partial^2)^{-1} &= q(\partial^2)^{-1}T^- - q^2\partial^1(\partial^2)^{-2}, \end{aligned} \tag{239}$$

$$\begin{aligned} \tau^{1/2}(\partial^1)^{-1} &= q^{-1}(\partial^1)^{-1}\tau^{1/2}, \\ \tau^{1/2}(\partial^2)^{-1} &= q(\partial^2)^{-1}\tau^{1/2}, \end{aligned} \tag{240}$$

$$\Lambda(\partial^i)^{-1} = q^{-2}(\partial^i)^{-1}\Lambda, \qquad i = 1, 2, \tag{241}$$

where Λ denotes a scaling operator.

As a next step we would like to find representations for the inverse partial derivatives. This can be achieved in the following way. In Ref. [38] it was shown that according to

$$\partial^i \triangleright F = \left(\partial^i_{\mathrm{cl}} + \partial^i_{\mathrm{cor}}\right) F \tag{242}$$

the representations of our partial derivatives can be divided up into a classical part and corrections vanishing in the undeformed limit $q \to 1$. Thus, seeking a solution to the equation

$$\partial^i \triangleright F = f \tag{243}$$

for given f it is reasonable to consider the following expression:

$$\begin{aligned} F = (\partial^i)^{-1} \triangleright f &= \frac{1}{\partial^i_{\mathrm{cl}} + \partial^i_{\mathrm{cor}}} f \\ &= \frac{1}{\partial^i_{\mathrm{cl}}\left(1 + (\partial^i_{\mathrm{cl}})^{-1}\partial^i_{\mathrm{cor}}\right)} f \\ &= \frac{1}{1 + (\partial^i_{\mathrm{cl}})^{-1}\partial^i_{\mathrm{cor}}} \cdot \frac{1}{\partial^i_{\mathrm{cl}}} f \\ &= \sum_{k=0}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1}\partial^i_{\mathrm{cor}}\right]^k (\partial^i_{\mathrm{cl}})^{-1} f. \end{aligned} \tag{244}$$

To apply this formula, we need to identify the contributions ∂^i_{cl} and ∂^i_{cor} the representations of our partial derivatives consist of. In the two-dimensional case, for example, we can read off from the expressions in (117) that

$$\begin{aligned}(\partial^1_{\text{cl}})f &= -q^{-1/2}D^2_{q^2}f(qx^1), \quad (\partial^1_{\text{cor}})f = 0,\\ (\partial^2_{\text{cl}})f &= q^{1/2}D^1_{q^2}f(q^2x^2), \quad (\partial^2_{\text{cor}})f = 0.\end{aligned} \tag{245}$$

Plugging this into formula (244) leaves us with

$$\begin{aligned}(\partial^1)^{-1} \triangleright f &= (\partial^1_{\text{cl}})^{-1} \triangleright f = -q^{1/2}(D^2_{q^2})^{-1}f(q^{-1}x^1),\\ (\partial^2)^{-1} \triangleright f &= (\partial^2_{\text{cl}})^{-1} \triangleright f = q^{-1/2}(D^1_{q^2})^{-1}f(q^{-2}x^2),\end{aligned} \tag{246}$$

where $(D^i_{q^a})^{-1}$ denotes the *Jackson integral* operator [61]. For $a > 0$, $q > 1$, and $x^i > 0$, the definition of the Jackson integral is given by

$$\begin{aligned}(D^i_{q^a})^{-1}\big|_0^{x^i} f &= -(1-q^a)\sum_{k=1}^{\infty}(q^{-ak}x^i)f(q^{-ak}x^i),\\ (D^i_{q^a})^{-1}\big|_{x^i}^{\infty} f &= -(1-q^a)\sum_{k=0}^{\infty}(q^{ak}x^i)f(q^{ak}x^i),\\ (D^i_{q^{-a}})^{-1}\big|_0^{x^i} f &= (1-q^{-a})\sum_{k=0}^{\infty}(q^{-ak}x^i)f(q^{-ak}x^i),\\ (D^i_{q^{-a}})^{-1}\big|_{x^i}^{\infty} f &= (1-q^{-a})\sum_{k=1}^{\infty}(q^{ak}x^i)f(q^{ak}x^i),\end{aligned} \tag{247}$$

and likewise for $a > 0$, $q > 1$, and $x^i < 0$,

$$\begin{aligned}(D^i_{q^a})^{-1}\big|_{x^i}^{0} f &= (1-q^a)\sum_{k=1}^{\infty}(q^{-ak}x^i)f(q^{-ak}x^i),\\ (D^i_{q^a})^{-1}\big|_{-\infty}^{x^i} f &= (1-q^a)\sum_{k=0}^{\infty}(q^{ak}x^i)f(q^{ak}x^i),\\ (D^i_{q^{-a}})^{-1}\big|_{x^i}^{0} f &= -(1-q^{-a})\sum_{k=0}^{\infty}(q^{-ak}x^i)f(q^{-ak}x^i),\end{aligned}$$

$$(D^i_{q^{-a}})^{-1}\big|_{-\infty}^{x^i} f = -(1-q^{-a})\sum_{k=1}^{\infty}(q^{ak}x^i)f(q^{ak}x^i). \tag{248}$$

From the discussion so far it is apparent that on an algebra of commutative functions the representations of inverse partial derivatives are given by expressions that depend on Jackson integrals and Jackson derivatives. But this is not the whole story, since the explicit form of a Jackson integral is determined by the choice of its limits. Realizing that the expression for $(\partial^i_{\rm cor})^{-1}$ is mainly given by a Jackson integral we can introduce definite integrals by

$$(\partial^i)^{-1}\big|_b^{\overline{x^i}} \triangleright f = \sum_{k=0}^{\infty}(-1)^k \left[(\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}}\partial^i_{\rm cor}\right]^k (\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}} f, \tag{249}$$

i.e. the lower and upper limits of all Jackson integrals appearing on the left side of Eq. (249) are respectively given by a constant b and the variable $x^{\overline{i}}$ (notice that the partial derivative ∂^i does not refer to the coordinate x^i but $x^{\overline{i}}$).

Now we would like to explore in which sense the operator in (249) is inverse to that in (242). To this end we perform the following calculations:

$$\begin{aligned}
&\partial^i \triangleright \left((\partial^i)^{-1}\big|_b^{\overline{x^i}} \triangleright f\right)\\
&= \sum_{k=0}^{\infty}(-1)^k\,(\partial^i_{\rm cl})\left[(\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}}\partial^i_{\rm cor}\right]^k (\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}} f\\
&\quad + \sum_{k=0}^{\infty}(-1)^k\,(\partial^i_{\rm cor})\left[\,(\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}}\,\partial^i_{\rm cor}\right]^k (\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}} f\\
&= \sum_{k=0}^{\infty}(-1)^k\left[\partial^i_{\rm cor}(\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}}\right]^k f\\
&\quad - \sum_{k=1}^{\infty}(-1)^k\left[\partial^i_{\rm cor}(\partial^i_{\rm cl})^{-1}\big|_b^{\overline{x^i}}\right]^k f\\
&= f,
\end{aligned} \tag{250}$$

and

$$(\partial^i)^{-1}\big|_b^{\overline{x^i}} \triangleright \left(\partial^i \triangleright f\right)$$

$$
\begin{aligned}
&= \sum_{k=0}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k (\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cl}} f \\
&\quad + \sum_{k=0}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k (\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} f \\
&= \sum_{k=0}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k \big(f \big|_b^{x^{\bar{i}}} \big) \\
&\quad - \sum_{k=1}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k f \\
&= \sum_{k=0}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k \big(f - f|_{x^{\bar{i}}=b} \big) \\
&\quad - \sum_{k=1}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k f \\
&= f - \sum_{k=0}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k \big(f|_{x^{\bar{i}}=b} \big). \qquad (251)
\end{aligned}
$$

In the above derivations we make use of the identities

$$
\partial^i_{\mathrm{cl}} (\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} f = f, \quad (\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cl}} f = f \big|_b^{x^{\bar{i}}}, \qquad (252)
$$

which are a direct consequence of the well-known relations

$$
\begin{aligned}
& D^i_{q^a} (D^i_{q^a})^{-1} \big|_b^{x^i} f = f, \\
& (D^i_{q^a})^{-1} \big|_b^{x^i} D^i_{q^a} f = f \big|_b^{x^i}. \qquad (253)
\end{aligned}
$$

We can now summarize that in contradiction to $\partial^i (\partial^i)^{-1} = (\partial^i)^{-1} \partial^i = 1$ we instead have

$$
\partial^i \triangleright \big((\partial^i)^{-1} \big|_b^{x^{\bar{i}}} \triangleright f \big) = f, \qquad (254)
$$

$$
(\partial^i)^{-1} \big|_b^{x^{\bar{i}}} \triangleright \big(\partial^i \triangleright f \big) = f - \hat{B}^i \big(f|_{x^{\bar{i}}=b} \big), \qquad (255)
$$

where, for brevity, we have introduced

$$\hat{B}^i \equiv \sum_{k=0}^{\infty} (-1)^k \left[(\partial^i_{\mathrm{cl}})^{-1} \big|_b^{x^{\bar{i}}} \partial^i_{\mathrm{cor}} \right]^k. \tag{256}$$

However, if the lower limit b tends to $-\infty$ and if we restrict attention to functions $f \in \widetilde{\mathcal{A}}$ with

$$\widetilde{\mathcal{A}} \equiv \Big\{ f \in \mathcal{A}, \quad \lim_{x^j \to -\infty} (x^j)^\alpha f = 0, \quad \text{for all } \alpha \in \mathbb{R} \text{ and } j \Big\}, \tag{257}$$

the representations (249) now respect all algebra relations for inverse partial derivatives [cf. (235) and (236)]. This observation motivates us to define the q-deformed integral over the interval $]-\infty, x^{\bar{i}}]$ as the mapping

$$\int_{-\infty}^{y^{\bar{i}}} d_L x^{\bar{i}} : \widetilde{\mathcal{A}} \longrightarrow \widetilde{\mathcal{A}}, \quad \int_{-\infty}^{y^{\bar{i}}} d_L x^{\bar{i}}\, f(x^j) \equiv \lim_{b \to -\infty} \big((\partial^i)^{-1} \big|_b^{x^{\bar{i}}} \triangleright f(x^j) \big). \tag{258}$$

If we want to deal with definite integrals over arbitrary intervals it is reasonable to generalize the definition of (258) by

$$\int_{z^{\bar{i}}}^{y^{\bar{i}}} d_L x^{\bar{i}} : \widetilde{\mathcal{A}} \longrightarrow \widetilde{\mathcal{A}} \otimes \widetilde{\mathcal{A}},$$

$$\int_{z^{\bar{i}}}^{y^{\bar{i}}} d_L x^{\bar{i}}\, f(x^j) \equiv \Big(\int_{-\infty}^{y^{\bar{i}}} d_L x^{\bar{i}}\, f(x^j) \Big) \otimes 1 - 1 \otimes \Big(\int_{-\infty}^{z^{\bar{i}}} d_L x^{\bar{i}}\, f(x^j) \Big). \tag{259}$$

We started our considerations for introducing q-deformed integrals from one specific representation of partial derivatives. But what we have done so far applies to the other representations for partial derivatives as well. In this manner, q-deformed integrals can be placed into four categories:

$$\begin{aligned}
\partial^i \triangleright f \quad &\Rightarrow \quad (\partial^i)^{-1} \big|_b^{x^{\bar{i}}} \triangleright f, \quad \int_{-\infty}^{y^{\bar{i}}} d_L x^{\bar{i}}\, f(x^j) \equiv \lim_{b \to -\infty} \big((\partial^i)^{-1} \big|_b^{x^{\bar{i}}} \triangleright f \big), \\
\hat{\partial}^i \bar{\triangleright} f \quad &\Rightarrow \quad (\hat{\partial}^i)^{-1} \big|_b^{x^{\bar{i}}} \bar{\triangleright} f, \quad \int_{-\infty}^{y^{\bar{i}}} d_{\bar{L}} x^{\bar{i}}\, f(x^j) \equiv \lim_{b \to -\infty} \big((\partial^i)^{-1} \big|_b^{x^{\bar{i}}} \bar{\triangleright} f \big),
\end{aligned} \tag{260}$$

$$f \triangleleft \hat{\partial}^i \quad \Rightarrow \quad f \triangleleft (\hat{\partial}^i)^{-1}\big|_b^{x^{\bar{i}}}, \quad \int_{-\infty}^{y^{\bar{i}}} d_R x^{\bar{i}}\, f(x^j) \equiv \lim_{b\to-\infty} \big(f \triangleleft (\hat{\partial}^i)^{-1}\big|_b^{x^{\bar{i}}}\big),$$
$$f \,\bar{\triangleleft}\, \partial^i \quad \Rightarrow \quad f \,\bar{\triangleleft}\, (\partial^i)^{-1}\big|_b^{x^{\bar{i}}}, \quad \int_{-\infty}^{y^{\bar{i}}} d_{\bar{R}} x^{\bar{i}}\, f(x^j) \equiv \lim_{b\to-\infty} \big(f \,\bar{\triangleleft}\, (\partial^i)^{-1}\big|_b^{x^{\bar{i}}}\big). \quad (261)$$

In complete analogy to (259) we have

$$\int_{z^{\bar{i}}}^{y^{\bar{i}}} d_{\bar{L}} x^{\bar{i}}\, f(x^j) \equiv \Big(\int_{-\infty}^{y^{\bar{i}}} d_{\bar{L}} x^{\bar{i}}\, f(x^j)\Big) \otimes 1 - 1 \otimes \Big(\int_{-\infty}^{z^{\bar{i}}} d_{\bar{L}} x^{\bar{i}}\, f(x^j)\Big), \quad (262)$$

but in contrast to this we now require for right integrals to hold

$$\int_{z^{\bar{i}}}^{y^{\bar{i}}} d_R x^{\bar{i}}\, f(x^j) \equiv 1 \otimes \Big(\int_{-\infty}^{y^{\bar{i}}} d_R x^{\bar{i}}\, f(x^j)\Big) - \Big(\int_{-\infty}^{z^{\bar{i}}} d_R x^{\bar{i}}\, f(x^j)\Big) \otimes 1,$$
$$\int_{z^{\bar{i}}}^{y^{\bar{i}}} d_{\bar{R}} x^{\bar{i}}\, f(x^j) \equiv 1 \otimes \Big(\int_{-\infty}^{y^{\bar{i}}} d_{\bar{R}} x^{\bar{i}}\, f(x^j)\Big) - \Big(\int_{-\infty}^{z^{\bar{i}}} d_{\bar{R}} x^{\bar{i}}\, f(x^j)\Big) \otimes 1. \quad (263)$$

It should be clear that the crossing-symmetries for representations of partial derivatives carry over to the corresponding integrals. For this reason, the relationship between the expressions for the various types of q-deformed integrals can be graphed as in Fig. 4. Applying these transition rules to the first relation in (246), we obtain as example

$$\begin{aligned} & (\partial^1)^{-1}\big|_b^{x^2} \triangleright f(x^1, x^2) = -q^{1/2} (D^2_{q^2})^{-1}\big|_b^{x^2} f(q^{-1}x^1, x^2) \\ \longleftrightarrow \quad & (\hat{\partial}^2)^{-1}\big|_b^{x^1} \,\bar{\triangleright}\, f(x^2, x^1) = -q^{-1/2} (D^1_{q^{-2}})^{-1}\big|_b^{x^1} f(qx^2, x^1), \\ \longleftrightarrow \quad & f(x^1, x^2) \,\bar{\triangleleft}\, (\partial^2)^{-1}\big|_b^{x^1} = -q^{1/2} (D^1_{q^2})^{-1}\big|_b^{x^1} f(x^1, q^{-1}x^2). \end{aligned} \quad (264)$$

5.2 Basic properties of q-deformed integrals

Next we would like to discuss some important properties of q-deformed integrals. First of all, the fact that the antipodes for the quantum space algebra intertwine the actions of partial derivatives [cf. (141) and (142)] also holds for the representations of inverse partial derivatives. This can be seen as follows:

$$\hat{\partial}^i \,\bar{\triangleright}\, \big(f(\ominus_R\, x^j)\big) = -\big(\partial^i \triangleright f\big)(\ominus_R\, x^j)$$




Figure 4: Crossing-symmetries for q-deformed integrals

$$
\begin{aligned}
&\Rightarrow \hat{\partial}^i \bar{\triangleright} \big[\big((\partial^i)^{-1} \triangleright f\big)(\ominus_R x^j)\big] = -\big(\partial^i \triangleright \big((\partial^i)^{-1} \triangleright f\big)\big)(\ominus_R x^j) \\
&\Rightarrow \hat{\partial}^i \bar{\triangleright} \big((\partial^i)^{-1} \triangleright f\big)(\ominus_R x^j) = -f(\ominus_R x^j) \\
&\Rightarrow (\hat{\partial}^i)^{-1} \bar{\triangleright} \big[\hat{\partial}^i \bar{\triangleright} \big[\big((\partial^i)^{-1} \triangleright f\big)(\ominus_R x^j)\big]\big] = -(\hat{\partial}^i)^{-1} \bar{\triangleright} \big(f(\ominus_R x^j)\big) \\
&\Rightarrow \big((\partial^i)^{-1} \triangleright f\big)(\ominus_R x^j) = -(\hat{\partial}^i)^{-1} \bar{\triangleright} \big(f(\ominus_R x^j)\big),
\end{aligned} \tag{265}
$$

where again the minus sign has to be dropped in the case of the Manin plane. In very much the same way, we obtain

$$
\big((\hat{\partial}^i)^{-1} \bar{\triangleright} f\big)(\ominus_{\bar{R}} x^j) = -(\partial^i)^{-1} \triangleright \big(f(\ominus_{\bar{R}} x^j)\big) \tag{266}
$$

and

$$
\begin{aligned}
\big(f \bar{\triangleleft} (\partial^i)^{-1}\big)(\ominus_{\bar{L}} x^j) &= -\big(f(\ominus_{\bar{L}} x^j)\big) \triangleleft (\hat{\partial}^i)^{-1}, \\
\big(f \triangleleft (\hat{\partial}^i)^{-1}\big)(\ominus_L x^j) &= -\big(f(\ominus_L x^j)\big) \bar{\triangleleft} (\partial^i)^{-1}.
\end{aligned} \tag{267}
$$

What we are doing next is to consider rules for integration by parts. To this end, let us first recall that the coproduct for partial derivatives leads to

a kind of q-deformed product rule, since we have

$$\begin{aligned}
\partial^i \triangleright (fg) &= (\partial^i \triangleright f)g + ((\mathcal{L}_\partial)^i_j \triangleright f)(\partial^j \triangleright g), \\
\hat{\partial}^i \bar{\triangleright} (fg) &= (\hat{\partial}^i \bar{\triangleright} f)g + ((\bar{\mathcal{L}}_\partial)^i_j \triangleright f)(\hat{\partial}^j \bar{\triangleright} g),
\end{aligned} \tag{268}$$

$$\begin{aligned}
(fg) \triangleleft \hat{\partial}^i &= f(g \triangleleft \hat{\partial}^i) + (f \triangleleft \hat{\partial}^j)(g \triangleleft (\mathcal{L}_\partial)^i_j), \\
(fg) \bar{\triangleleft} \partial^i &= f(g \bar{\triangleleft} \partial^i) + (f \bar{\triangleleft} \partial^j)(g \triangleleft (\bar{\mathcal{L}}_\partial)^i_j).
\end{aligned} \tag{269}$$

Hitting the above equations with the corresponding integral operator and rearranging terms, we get

$$\begin{aligned}
\int_{-\infty}^{y^{\bar{i}}} d_L x^{\bar{i}} \, (\partial^i \triangleright f)g &= fg|_{-\infty}^{y^{\bar{i}}} - \int_{-\infty}^{y^{\bar{i}}} d_L x^{\bar{i}} \, ((\mathcal{L}_\partial)^i_j \triangleright f)(\partial^j \triangleright g), \\
\int_{-\infty}^{y^{\bar{i}}} d_{\bar{L}} x^{\bar{i}} \, (\hat{\partial}^i \bar{\triangleright} f)g &= fg|_{-\infty}^{y^{\bar{i}}} - \int_{-\infty}^{y^{\bar{i}}} d_{\bar{L}} x^{\bar{i}} \, ((\bar{\mathcal{L}}_\partial)^i_j \triangleright f)(\hat{\partial}^j \bar{\triangleright} g),
\end{aligned} \tag{270}$$

$$\begin{aligned}
\int_{-\infty}^{y^{\bar{i}}} d_R x^{\bar{i}} \, f(g \triangleleft \hat{\partial}^i) &= fg|_{-\infty}^{y^{\bar{i}}} - \int_{-\infty}^{y^{\bar{i}}} d_R x^{\bar{i}} \, (f \triangleleft \hat{\partial}^j)(g \triangleleft (\mathcal{L}_\partial)^i_j), \\
\int_{-\infty}^{y^{\bar{i}}} d_{\bar{R}} x^{\bar{i}} \, f(g \bar{\triangleleft} \partial^i) &= fg|_{-\infty}^{y^{\bar{i}}} - \int_{-\infty}^{y^{\bar{i}}} d_{\bar{R}} x^{\bar{i}} \, (f \bar{\triangleleft} \partial^j)(g \triangleleft (\bar{\mathcal{L}}_\partial)^i_j).
\end{aligned} \tag{271}$$

It is not very difficult to convince oneself that the above identities also hold for integrals over arbitrary intervals. In the case of the two-dimensional quantum plane the rules in (270) and (271) explicitly read as

$$\begin{aligned}
&\int_{-\infty}^{y^2} d_L x^2 \, (\partial^1 \triangleright f) \circledast g = \\
&\qquad f \circledast g|_{-\infty}^{y^2} - \int_{-\infty}^{y^2} d_L x^2 \, (\Lambda^{-3/4} \tau^{-1/4} \triangleright f) \circledast (\partial^1 \triangleright g),
\end{aligned} \tag{272}$$

$$\begin{aligned}
&\int_{-\infty}^{y^1} d_L x^1 \, (\partial^2 \triangleright f) \circledast g = \\
&\qquad f \circledast g|_{-\infty}^{y^1} - \int_{-\infty}^{y^1} d_L x^1 \, (\Lambda^{-3/4} \tau^{1/4} \triangleright f) \circledast (\partial^2 \triangleright g)
\end{aligned}$$

$$+ q\lambda \int_{-\infty}^{y^1} d_L x^1 \, (\Lambda^{-3/4}\tau^{-1/4}T^+ \triangleright f) \circledast (\partial^1 \triangleright g), \tag{273}$$

$$\int_{-\infty}^{y^2} d_{\bar{L}} x^2 \, (\hat{\partial}^1 \bar{\triangleright} f) \circledast g =$$
$$f \circledast g|_{-\infty}^{y^2} - \int_{-\infty}^{y^2} d_{\bar{L}} x^2 \, (\Lambda^{3/4}\tau^{1/4} \triangleright f) \circledast (\hat{\partial}^1 \bar{\triangleright} g)$$
$$- q^{-1}\lambda \int_{-\infty}^{y^2} d_{\bar{L}} x^2 \, (\Lambda^{3/4}\tau^{-1/4}T^- \triangleright f) \circledast (\hat{\partial}^2 \bar{\triangleright} g), \tag{274}$$

$$\int_{-\infty}^{y^1} d_{\bar{L}} x^1 \, (\hat{\partial}^2 \bar{\triangleright} f) g =$$
$$f \circledast g|_{-\infty}^{y^1} - \int_{-\infty}^{y^1} d_{\bar{L}} x^1 \, (\Lambda^{3/4}\tau^{-1/4} \triangleright f) \circledast (\hat{\partial}^2 \bar{\triangleright} g), \tag{275}$$

and likewise

$$\int_{-\infty}^{y^2} d_R x^2 \, f \circledast (g \triangleleft \hat{\partial}^1) =$$
$$f \circledast g|_{-\infty}^{y^2} - \int_{-\infty}^{y^2} d_R x^2 \, (f \triangleleft \hat{\partial}^1) \circledast (g \triangleleft \Lambda^{-3/4}\tau^{-1/4}), \tag{276}$$

$$\int_{-\infty}^{y^1} d_R x^1 \, f \circledast (g \triangleleft \hat{\partial}^2) =$$
$$f \circledast g|_{-\infty}^{y^1} - \int_{-\infty}^{y^1} d_R x^1 \, (f \triangleleft \hat{\partial}^2) \circledast (g \triangleleft \Lambda^{-3/4}\tau^{1/4})$$
$$+ q\lambda \int_{-\infty}^{y^1} d_R x^1 \, (f \triangleleft \hat{\partial}^1) \circledast (g \triangleleft \Lambda^{-3/4}\tau^{-1/4}T^+), \tag{277}$$

$$\int_{-\infty}^{y^2} d_{\bar{R}} x^2 \, f \circledast (g \bar{\triangleleft} \partial^1) =$$
$$f \circledast g|_{-\infty}^{y^2} - \int_{-\infty}^{y^2} d_{\bar{R}} x^2 \, (f \bar{\triangleleft} \partial^1) \circledast (g \triangleleft \Lambda^{3/4}\tau^{1/4})$$

$$- q^{-1}\lambda \int_{-\infty}^{y^2} d_{\bar{R}}x^2\, (f \,\bar{\triangleleft}\, \partial^2) \circledast (g \triangleleft \Lambda^{3/4}\tau^{-1/4}T^-), \tag{278}$$

$$\int_{-\infty}^{y^1} d_{\bar{R}}x^1\, f \circledast (g \,\bar{\triangleleft}\, \partial^2) =$$

$$f \circledast g|_{-\infty}^{y^1} - \int_{-\infty}^{y^1} d_{\bar{R}}x^1\, (f \,\bar{\triangleleft}\, \partial^2) \circledast (g \triangleleft \Lambda^{3/4}\tau^{-1/4}). \tag{279}$$

One readily checks the above identities by inserting the expressions of (18) and (19) into (270) and (271).

Now, we come to another important point concerning translations of integrals. To this end, we take up the relations in (186)-(189) again and proceed as follows:

$$\partial^i \overset{x}{\triangleright} \exp(x^k|\partial^l)_{\bar{R},L} = \exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial}{\circledast} \partial^i$$

$$\Rightarrow \quad \partial^i \overset{x}{\triangleright} \exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial|y}{\triangleright} \int_{-\infty}^{y^{\bar{i}}} d_L y^{\bar{i}}\, f(y^j) =$$

$$\big(\exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial}{\circledast} \partial^i\big) \overset{\partial|y}{\triangleright} \int_{-\infty}^{y^{\bar{i}}} d_L \tilde{y}^{\bar{i}}\, f(\tilde{y}^j)$$

$$\Rightarrow \quad \int_{-\infty}^{x^{\bar{i}}} d_L \tilde{x}^{\bar{i}}\, \tilde{\partial}^i \overset{\tilde{\partial}|\tilde{x}}{\triangleright} \exp(\tilde{x}^k|\partial^l)_{\bar{R},L} \overset{\partial|y}{\triangleright} \int_{-\infty}^{y^{\bar{i}}} d_L \tilde{y}^{\bar{i}}\, f(\tilde{y}^j) =$$

$$\int_{-\infty}^{x^{\bar{i}}} d_L \tilde{x}^{\bar{i}}\, \exp(\tilde{x}^k|\partial^l)_{\bar{R},L} \overset{\partial|y}{\triangleright} f(y^j)$$

$$\Rightarrow \quad \exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial|y}{\triangleright} \int_{-\infty}^{y^{\bar{i}}} d_L \tilde{y}^{\bar{i}}\, f(\tilde{y}^j) = \int_{-\infty}^{x^{\bar{i}}} d_L \tilde{x}^{\bar{i}}\, \exp(\tilde{x}^k|\partial^l)_{\bar{R},L} \overset{\partial|y}{\triangleright} f(y^j)$$

$$\Rightarrow \quad \int_{-\infty}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{y}^{\overline{i}} \, f(\tilde{y}^j) = \int_{-\infty}^{x^{\overline{i}}} d_L \tilde{x}^{\overline{i}} \, f(\tilde{x}^k \oplus_{\bar{L}} y^j). \tag{280}$$

In the above derivation we suppose $f \in \widetilde{\mathcal{A}}$ and apply (255). For the last step we make use of (214).

If we consider integrals over finite intervals, we have to work a little bit harder. First of all, it is necessary to realize that

$$\begin{aligned} \partial^i \overset{y,z}{\triangleright} \int_{z^{\overline{i}}}^{y^{\overline{i}}} d_L \tilde{y}^{\overline{i}} \, f(\tilde{y}^k) &= \partial^i \overset{y}{\triangleright} \int_{-\infty}^{y^{\overline{i}}} d_L \tilde{y}^{\overline{i}} \, f(\tilde{y}^k) \otimes 1 - 1 \otimes \partial^i \overset{z}{\triangleright} \int_{-\infty}^{z^{\overline{i}}} d_L \tilde{z}^{\overline{i}} \, f(\tilde{z}^l) \\ &= f(y^k) \otimes 1 - 1 \otimes f(z^l), \end{aligned} \tag{281}$$

and analogously

$$\begin{aligned} & \exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial|y,z}{\triangleright} \int_{z^{\overline{i}}}^{y^{\overline{i}}} d_L \tilde{y}^{\overline{i}} \, f(\tilde{y}^j) \\ &= \exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial|y}{\triangleright} \int_{-\infty}^{y^{\overline{i}}} d_L \tilde{y}^{\overline{i}} \, f(\tilde{y}^j) \otimes 1 - 1 \otimes \exp(x^k|\partial^l)_{\bar{R},L} \overset{\partial|z}{\triangleright} \int_{-\infty}^{z^{\overline{i}}} d_L \tilde{z}^{\overline{i}} \, f(\bar{z}^j) \\ &= \int_{-\infty}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{y}^{\overline{i}} \, f(\tilde{y}^j) \otimes 1 - 1 \otimes \int_{-\infty}^{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}} d_L \tilde{z}^{\overline{i}} \, f(\tilde{z}^j) \\ &= \int_{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{y}^{\overline{i}} \, f(\tilde{y}^j). \end{aligned} \tag{282}$$

With these identities at hand the considerations in (280) now lead to

$$\int_{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} \, f(\tilde{x}^k) = \int_{-\infty}^{x^{\overline{i}}} d_L \tilde{x}^{\overline{i}} \, f(\tilde{x}^k \oplus_{\bar{L}} y^j) - \int_{-\infty}^{x^{\overline{i}}} d_L \tilde{x}^{\overline{i}} \, f(\tilde{x}^k \oplus_{\bar{L}} z^j)$$

$$
\begin{aligned}
&= \int_{-\infty}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) - \int_{-\infty}^{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) \\
&= \int_{-\infty}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) - \int_{-\infty}^{x^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) \\
&- \int_{-\infty}^{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) + \int_{-\infty}^{x^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) \\
&= \int_{x^{\overline{i}}}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) - \int_{x^{\overline{i}}}^{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k), \qquad (283)
\end{aligned}
$$

being equivalent to

$$
\int_{x^{\overline{i}}}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) = \int_{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}}^{(x^k \oplus_{\bar{L}} y^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k) + \int_{x^{\overline{i}}}^{(x^k \oplus_{\bar{L}} z^j)^{\overline{i}}} d_L \tilde{x}^{\overline{i}} f(\tilde{x}^k). \qquad (284)
$$

Everything so far applies equally well to the other types of integrals, i.e. we additionally have

$$
\begin{aligned}
\int_{-\infty}^{(x^k \oplus_L y^j)^{\overline{i}}} d_{\bar{L}} \tilde{x}^{\overline{i}} f(\tilde{x}^j) &= \int_{-\infty}^{x^{\overline{i}}} d_{\bar{L}} \tilde{x}^{\overline{i}} f(\tilde{x}^k \oplus_L y^j) \\
\int_{-\infty}^{(y^j \oplus_{\bar{R}} x^k)^{\overline{i}}} d_R \tilde{x}^{\overline{i}} f(\tilde{x}^j) &= \int_{-\infty}^{x^{\overline{i}}} d_R \tilde{x}^{\overline{i}} f(y^j \oplus_{\bar{R}} \tilde{x}^k), \\
\int_{-\infty}^{(y^j \oplus_R x^k)^{\overline{i}}} d_{\bar{R}} \tilde{x}^{\overline{i}} f(\tilde{x}^j) &= \int_{-\infty}^{x^{\overline{i}}} d_{\bar{R}} \tilde{x}^{\overline{i}} f(y^j \oplus_R \tilde{x}^k), \qquad (285)
\end{aligned}
$$

and

$$\int_{x^{\bar{i}}}^{(x^k\oplus_L y^j)^{\bar{i}}} d_{\bar{L}}\tilde{x}^{\bar{i}}\, f(\tilde{x}^k) = \int_{(x^k\oplus_L z^j)^{\bar{i}}}^{(x^k\oplus_L y^j)^{\bar{i}}} d_{\bar{L}}\tilde{x}^{\bar{i}}\, f(\tilde{x}^k) + \int_{x^{\bar{i}}}^{(x^k\oplus_L z^j)^{\bar{i}}} d_{\bar{L}}\tilde{x}^{\bar{i}}\, f(\tilde{x}^k),$$

$$\int_{x^{\bar{i}}}^{(y^j\oplus_{\bar{R}} x^k)^{\bar{i}}} d_R\tilde{x}^{\bar{i}}\, f(\tilde{x}^k) = \int_{(z^j\oplus_{\bar{R}} x^k)^{\bar{i}}}^{(y^j\oplus_{\bar{R}} x^k)^{\bar{i}}} d_R\tilde{x}^{\bar{i}}\, f(\tilde{x}^k) + \int_{x^{\bar{i}}}^{(z^j\oplus_{\bar{R}} x^k)^{\bar{i}}} d_R\tilde{x}^{\bar{i}}\, f(\tilde{x}^k),$$

$$\int_{x^{\bar{i}}}^{(y^j\oplus_R x^k)^{\bar{i}}} d_{\bar{R}}\tilde{x}^{\bar{i}}\, f(\tilde{x}^k) = \int_{(z^j\oplus_R x^k)^{\bar{i}}}^{(y^j\oplus_R x^k)^{\bar{i}}} d_{\bar{R}}\tilde{x}^{\bar{i}}\, f(\tilde{x}^k) + \int_{x^{\bar{i}}}^{(z^j\oplus_R x^k)^{\bar{i}}} d_{\bar{R}}\tilde{x}^{\bar{i}}\, f(\tilde{x}^k). \qquad (286)$$

For a correct understanding of the relations in (285) one should keep in mind that the set $\widetilde{\mathcal{A}}$ [cf. Eq. (257)] is invariant under translations, i.e. it holds

$$f \in \widetilde{\mathcal{A}} \quad \Rightarrow \quad f(x^i \oplus_{L/\bar{L}} y^j) \in \widetilde{\mathcal{A}}, \quad f(\ominus_{L/\bar{L}} x^i) \in \widetilde{\mathcal{A}}. \qquad (287)$$

In other words, translations do not change the boundary conditions a function shows at infinity. This assertion follows from a direct inspection of the operator expressions for q-deformed translations (see for example Ref. [41]).

5.3 Integrals over the whole space and their translation invariance

Up to now we have considered one-dimensional integrals only. We are now looking for integrals over the entire space. In the work of Ref. [39] such integrals were obtained by applying one-dimensional integrals in succession, i.e.

$$\int_{-\infty}^{+\infty} d_L^n x\, f \equiv \int_{-\infty}^{+\infty} d_L x^{\bar{n}} \ldots \int_{-\infty}^{+\infty} d_L x^{\bar{2}} \int_{-\infty}^{+\infty} d_L x^{\bar{1}}\, f(x^i)$$
$$= \lim_{x^{\bar{1}},\ldots,x^{\bar{n}}\to\infty} \left((\partial^n)^{-1}\Big|_{-\infty}^{x^{\bar{n}}} \ldots (\partial^1)^{-1}\Big|_{-\infty}^{x^{\bar{1}}} \triangleright f\right),$$

$$\int_{-\infty}^{+\infty} d_R^n x\, f \equiv \int_{-\infty}^{+\infty} d_R x^1 \int_{-\infty}^{+\infty} d_R x^2 \ldots \int_{-\infty}^{+\infty} d_R x^n\, f(x^i)$$
$$= \lim_{x^1,\ldots,x^n\to\infty} \big(f \triangleleft (\hat{\partial}^{\bar{1}})^{-1}\big|_{-\infty}^{x^1} \ldots (\hat{\partial}^{\bar{n}})^{-1}\big|_{-\infty}^{x^n} \big). \qquad (288)$$

There seems to be one difficulty with these definitions, since they can depend on the order in which the one-dimensional integrals are performed. However, in the above definitions we take for all the lower and upper limits of integration $-\infty$ and $+\infty$, respectively. For this reason, the one-dimensional integrals being of the form (244) contribute to the expressions in (288) by their summands with $k = 0$, only. This observation relies on the fact that all summands with $k > 0$ lead to surface terms that can be neglected. This way, the expressions in (288) should simplify to

$$\int_{-\infty}^{+\infty} d_L^n x\, f(x^i) = (\partial_{\mathrm{cl}}^n)^{-1}\big|_{-\infty}^{\infty} \ldots (\partial_{\mathrm{cl}}^1)^{-1}\big|_{-\infty}^{\infty} \triangleright f,$$
$$\int_{-\infty}^{+\infty} d_R^n x\, f(x^i) = f \triangleleft (\hat{\partial}_{\mathrm{cl}}^{\bar{1}})^{-1}\big|_{-\infty}^{\infty} \ldots (\hat{\partial}_{\mathrm{cl}}^{\bar{n}})^{-1}\big|_{-\infty}^{\infty}. \qquad (289)$$

The point now is that the integral operators $(\partial_{\mathrm{cl}}^i)^{-1}$, $i = 1, \ldots, n$, commute among each other up to certain powers of q, as can be seen from a direct inspection of their explicit form [cf. (246)]. Thus, we can conclude that the above definitions do not really depend on the order of the one-dimensional integrals, since changing their order leads to expressions that differ from the original ones by a global factor, only.

Next, we wish to find the explicit form of the integrals over the whole space for the cases we are interested in for physical reasons. This can be done as sketched above. First, we identify the expressions for each ∂_{cl}^i. Taking the corresponding inverse and plugging it into the formulae of (289), we finally obtain

(i) (quantum plane)

$$\int_{-\infty}^{+\infty} d_L^2 x\, f(x^1, x^2) \equiv -(D_{q^2}^1)^{-1}\big|_{-\infty}^{\infty} q^{-2\hat{n}_{x^2}} (D_{q^2}^2)^{-1}\big|_{-\infty}^{\infty} q^{-\hat{n}_{x^1}} f, \qquad (290)$$

(ii) (three-dimensional Euclidean space)

$$\int_{-\infty}^{+\infty} d_L^3 x\, f(x^+, x^3, x^-) \equiv$$
$$(D^+_{q^4})^{-1}\big|^{\infty}_{-\infty}(D^3_{q^2})^{-1}\big|^{\infty}_{-\infty} q^{2\hat{n}_{x^+}}(D^-_{q^4})^{-1}\big|^{\infty}_{-\infty} q^{2\hat{n}_{x^3}} f, \tag{291}$$

(iii) (four-dimensional Euclidean space)

$$\int_{-\infty}^{+\infty} d_L^4 x\, f(x^4, x^3, x^2, x^1) \equiv$$
$$(D^1_{q^2})^{-1}\big|^{\infty}_{-\infty} q^{\hat{n}_{x^2}+\hat{n}_{x^3}}(D^2_{q^2})^{-1} q^{\hat{n}_4}\big|^{\infty}_{-\infty}(D^3_{q^2})^{-1}\big|^{\infty}_{-\infty} q^{\hat{n}_{x^4}}(D^4_{q^2})^{-1}\big|^{\infty}_{-\infty} f, \tag{292}$$

(iv) (q-deformed Minkowski space)

$$\int_{-\infty}^{+\infty} d_L^4 x\, f(r^2, x^-, x^{3/0}, x^+) \equiv$$
$$-q\lambda_+^{-3}(D^{r^2}_{q^{-2}})^{-1}\big|^{\infty}_{-\infty} q^{2(\hat{n}_{x^+}+\hat{n}_{x^{3/0}}+\hat{n}_{x^-})}(D^+_{q^{-2}})^{-1}\big|^{\infty}_{-\infty}$$
$$\times(D^{3/0}_{q^{-2}})^{-1}\big|^{\infty}_{-\infty} q^{2\hat{n}_{r^2}}(D^-_{q^{-2}})^{-1}\big|^{\infty}_{-\infty} q^{2\hat{n}_{r^2}} f. \tag{293}$$

Notice that we have added Appendix B to explain in more detail the reasonings leading to the expression in (293).

Obviously, the above considerations apply to all types of one-dimensional integrals without any difficulties. As we know, it suffices to concern ourselves with one type of integrals, since the expressions for the other versions are obtained most easily by the crossing symmetries graphed in Fig. 4. Thus, we concretely have the correspondences

$$\int_{-\infty}^{+\infty} d_L^n x\, f(x^i) = (\partial^n)^{-1}\big|^{\infty}_{-\infty} \ldots (\partial^1)^{-1}\big|^{\infty}_{-\infty} \triangleright f$$

$$\overset{\substack{i\leftrightarrow\bar{i}\\ q\leftrightarrow 1/q}}{\longleftrightarrow} \quad \int_{-\infty}^{+\infty} d^n_{\bar{L}}x\, \tilde{f}(x^i) = (\hat{\partial}^{\bar{n}})^{-1}\big|_{-\infty}^{\infty} \ldots (\hat{\partial}^{\bar{1}})^{-1}\big|_{-\infty}^{\infty} \bar{\triangleright}\, \tilde{f}, \tag{294}$$

$$\int_{-\infty}^{+\infty} d^n_{\bar{R}}x\, f(x^i) = f \,\bar{\triangleleft}\, (\partial^1)^{-1}\big|_{-\infty}^{\infty} \ldots (\partial^n)^{-1}\big|_{-\infty}^{\infty}$$

$$\overset{\substack{i\leftrightarrow\bar{i}\\ q\leftrightarrow 1/q}}{\longleftrightarrow} \quad \int_{-\infty}^{+\infty} d^n_{R}x\, \tilde{f}(x^i) = \tilde{f} \triangleleft (\hat{\partial}^{\bar{1}})^{-1}\big|_{-\infty}^{\infty} \ldots (\hat{\partial}^{\bar{n}})^{-1}\big|_{-\infty}^{\infty}, \tag{295}$$

and

$$\int_{-\infty}^{+\infty} d^n_{R}x\, \tilde{f}(x^i) \overset{i\leftrightarrow\bar{i}}{\longleftrightarrow} (-1)^n \int_{-\infty}^{+\infty} d^n_{\bar{L}}x\, \tilde{f}(x^i), \tag{296}$$

$$\int_{-\infty}^{+\infty} d^n_{\bar{R}}x\, f(x^i) \overset{i\leftrightarrow\bar{i}}{\longleftrightarrow} (-1)^n \int_{-\infty}^{+\infty} d^n_{L}x\, f(x^i),$$

where $\overset{\substack{i\leftrightarrow\bar{i}\\ q\leftrightarrow 1/q}}{\longleftrightarrow}$ and $\overset{i\leftrightarrow\bar{i}}{\longleftrightarrow}$ respectively indicate transitions via the substitutions

$$(D^i_{q^a})^{-1} \to (D^{\bar{i}}_{q^{-a}})^{-1}, \quad \hat{n}_{x^i} \to \hat{n}_{x^{\bar{i}}}, \quad x^i \to x^{\bar{i}}, \quad q \to q^{-1}, \tag{297}$$

and

$$(D^i_{q^a})^{-1} \to (D^{\bar{i}}_{q^a})^{-1}, \quad \hat{n}_{x^i} \to \hat{n}_{x^{\bar{i}}}, \quad x^i \to x^{\bar{i}}. \tag{298}$$

Furthermore, one should notice that the tilde on the symbol for the function shall remind us of the fact that the corresponding expression refers to reversed normal ordering. We would like to illustrate the transition rules in (294)-(296) by the following example:

$$\int_{-\infty}^{+\infty} d^2_{L}x\, f(x^1,x^2) \equiv -(D^1_{q^2})^{-1}\big|_{-\infty}^{\infty} q^{-2\hat{n}_{x^2}} (D^2_{q^2})^{-1}\big|_{-\infty}^{\infty} q^{-\hat{n}_{x^1}} f$$

$$\overset{\substack{i\leftrightarrow\bar{i}\\ q\leftrightarrow 1/q}}{\longleftrightarrow} \quad \int_{-\infty}^{+\infty} d^2_{\bar{L}}x\, f(x^2,x^1) = -(D^2_{q^{-2}})^{-1}\big|_{-\infty}^{\infty} q^{2\hat{n}_{x^1}} (D^1_{q^{-2}})^{-1}\big|_{-\infty}^{\infty} q^{\hat{n}_{x^2}} f$$

$$\overset{i\leftrightarrow\bar{i}}{\longleftrightarrow} \quad \int_{-\infty}^{+\infty} d_R^2 x\, f(x^2,x^1) = -(D^1_{q^{-2}})^{-1}\big|_{-\infty}^{\infty} q^{2\hat{n}_{x^2}} (D^2_{q^{-2}})^{-1}\big|_{-\infty}^{\infty} q^{\hat{n}_{x^1}} f. \tag{299}$$

Remarkably, q-deformed integrals over the whole space can also be obtained in a different manner, i.e. by seeking expressions being translationally invariant. This way we begin by defining an integral over the whole space as a mapping

$$\int : \widetilde{\mathcal{A}} \to \mathbb{C}, \tag{300}$$

being left and right invariant in the sense

$$\int \partial^i \triangleright f = \int f \triangleleft \partial^i = \varepsilon(\partial^i) \int f = 0. \tag{301}$$

Notice that this kind of invariance also implies invariance under the action of generators of the quantum algebra describing the underlying symmetry, i.e.

$$\int L^{ij} \triangleright f = \int f \triangleleft L^{ij} = \varepsilon(L^{ij}) \int f = 0. \tag{302}$$

This can be seen rather easily from the fact that the generators of the quantum algebra are realized within the differential calculus by

$$L^{ij} \equiv k(P_A)^{ij}{}_{kl}\, \Lambda^\alpha X^k \partial^l = k'(P_A)^{ij}{}_{kl}\, \partial^k X^l \Lambda^\alpha, \tag{303}$$

where k and k' stand for suitable constants, while Λ^α and P_A denote a scaling operator and a q-analog of an antisymmetrizer, respectively.

That the integrals given by (288) indeed satisfy (301) can be readily checked by using the commutation relations involving inverse partial derivatives [see for example (255)]. In this respect, we should realize that each time we commute a partial derivative with an inverse one a one-dimensional integral disappears and surface terms, which vanish at infinity, arise. In the case of the two-dimensional quantum plane, for example, we have

$$\begin{aligned} &\lim_{x^1,x^2\to\infty} (\partial^2)^{-1}\big|_{-\infty}^{x^1} (\partial^1)^{-1}\big|_{-\infty}^{x^2} \triangleright (\partial^1 \triangleright f) \\ &= \lim_{x^1,x^2\to\infty} (\partial^2)^{-1}\big|_{-\infty}^{x^1} \triangleright f = 0, \end{aligned} \tag{304}$$

$$\lim_{x^1,x^2\to\infty}(\partial^2)^{-1}\big|_{-\infty}^{x^1}(\partial^1)^{-1}\big|_{-\infty}^{x^2}\triangleright(\partial^2\triangleright f)$$
$$= q^{-1}\lim_{x^1,x^2\to\infty}(\partial^1)^{-1}\big|_{-\infty}^{x^2}\triangleright f = 0, \tag{305}$$

where it is assumed that $f\in\widetilde{\mathcal{A}}$. It should be clear that these reasonings require for the representations of the ∂^i to be of the same type as the representations of the $(\partial^i)^{-1}$. However, from (146) and (147) we know that the antipode and its inverse enable us to transform left and right actions into each other. In this manner, we can proceed as follows:

$$\lim_{x^{\bar{1}},\ldots,x^{\bar{n}}\to\infty}\big[(\partial^n)^{-1}\big|_{-\infty}^{x^{\bar{n}}}\ldots(\partial^1)^{-1}\big|_{-\infty}^{x^{\bar{1}}}\triangleright(f\triangleleft\hat{\partial}^i)\big]$$
$$= \lim_{x^{\bar{1}},\ldots,x^{\bar{n}}\to\infty}\big[(\partial^n)^{-1}\big|_{-\infty}^{x^{\bar{n}}}\ldots(\partial^1)^{-1}\big|_{-\infty}^{x^{\bar{1}}}\triangleright(-\hat{\partial}^j S^{-1}(\mathcal{L}_\partial)^i_j\triangleright f)\big]$$
$$= 0. \tag{306}$$

The above considerations also apply for right integrals. Thus, we have

$$\lim_{x^1,\ldots,x^n\to\infty}\big[(\partial^i\triangleright f)\triangleleft(\hat{\partial}^{\bar{1}})^{-1}\big|_{-\infty}^{x^1}\ldots(\hat{\partial}^{\bar{n}})^{-1}\big|_{-\infty}^{x^n}\big]$$
$$= \lim_{x^1,\ldots,x^n\to\infty}\big[(f\triangleleft\hat{\partial}^i)\triangleleft(\hat{\partial}^{\bar{1}})^{-1}\big|_{-\infty}^{x^1}\ldots(\hat{\partial}^{\bar{n}})^{-1}\big|_{-\infty}^{x^n}\big]$$
$$= 0. \tag{307}$$

Moreover, conjugate actions of partial derivatives can be expressed by unconjugate ones and vice versa, since we have [18, 71]

$$\hat{\partial}^i\,\bar{\triangleright}\,f = k_1\partial^i\triangleright(\Lambda\triangleright f) + k_2(g_{ij}\partial^i\partial^i)\triangleright(\Lambda\triangleright(x^i\circledast f)),$$
$$\partial^i\triangleright f = \tilde{k}_1\hat{\partial}^i\,\bar{\triangleright}\,(\tilde{\Lambda}\triangleright f) + \tilde{k}_2(g_{ij}\hat{\partial}^i\hat{\partial}^i)\,\bar{\triangleright}\,(\tilde{\Lambda}\triangleright(x^i\circledast f)), \tag{308}$$

and

$$f\triangleleft\hat{\partial}^i = k_3(f\triangleleft\Lambda^{-1})\,\bar{\triangleleft}\,\partial^i + k_4((f\circledast x^i)\triangleleft\Lambda^{-1})\,\bar{\triangleleft}\,(\partial^i\partial^i g_{ij}),$$
$$f\,\bar{\triangleleft}\,\partial^i = \tilde{k}_3(f\triangleleft\tilde{\Lambda}^{-1})\,\bar{\triangleleft}\,\hat{\partial}^i + \tilde{k}_4((f\circledast x^i)\triangleleft\tilde{\Lambda}^{-1})\triangleleft(\hat{\partial}^i\hat{\partial}^i g_{ij}), \tag{309}$$

with suitable constants $k_i,\tilde{k}_i\in\mathbb{R}$ and scaling operators Λ and $\tilde{\Lambda}$. Together

with (306) and (307) these relations give rise to the identities

$$\begin{aligned}\int d^n_A x(\partial^i \triangleright f) &= \int d^n_A x(\hat{\partial}^i \bar{\triangleright} f)\\ = \int d^n_A x(f \triangleleft \hat{\partial}^i) &= \int d^n_A x(f \bar{\triangleleft} \partial^i) = 0,\end{aligned} \tag{310}$$

where $A \in \{L, \bar{L}, R, \bar{R}\}$.

Translation invariance of integrals can help to simplify the rules for integration by parts. This becomes quite clear from the calculation

$$\begin{aligned}\int_{-\infty}^{+\infty} d_{L/R}x^n\, f \circledast (\partial^i \triangleright g) &= \int_{-\infty}^{+\infty} d_{L/R}x^n\, (f \triangleleft \partial^i) \circledast g\\ &\quad + \int_{-\infty}^{+\infty} d_{L/R}x^n\, (\mathcal{L}_\partial)^i_j \triangleright \big[(f \triangleleft \partial^j) \circledast g\big]\\ &= \int_{-\infty}^{+\infty} d_{L/R}x^n\, (f \triangleleft \partial^i) \circledast g.\end{aligned} \tag{311}$$

The first step is an application of the coproduct for partial derivatives. For the second step we have to recall that the entries of the L-matrix are elements of the symmetry algebra. [To be more precise, we should mention that the entries of the L-matrix depend on a scaling operator. But the invariance of our integrals under the action of this scaling operator is a rather subtle point. See the discussion of (324)]. Applying (302) together with $\varepsilon(\mathcal{L}^i_j) = \delta^i_j$ then yields the final result. By a slight modification we can repeat these arguments for the symmetry generators L^{ij}. In doing so, we get

$$\begin{aligned}&\int_{-\infty}^{+\infty} d_{L/R}x^n\, f \circledast (L^{ij} \triangleright g)\\ &= \int_{-\infty}^{+\infty} d_{L/R}x^n\, (L^{ij})_{(2)} \triangleright \big(\big[S^{-1}((L^{ij})_{(1)}) \triangleright f\big] \circledast g\big)\end{aligned}$$

$$= \int_{-\infty}^{+\infty} d_{L/R}x^n\, \varepsilon((L^{ij})_{(2)})\, \left[\left(f \triangleleft (L^{ij})_{(1)}\right) \circledast g\right]$$
$$= \int_{-\infty}^{+\infty} d_{L/R}x^n\, \left(f \triangleleft L^{ij}\right) \circledast g, \tag{312}$$

where the second step again makes use of (302).

Next, we would like to rewrite the condition of translation invariance in terms of q-translations. To this end, we first reformulate the properties in (310) by inserting the expressions for the coregular actions [cf. (156) and (157)]. The non-degeneracy of the pairings then implies

$$\left\langle \partial^i, f_{(\bar{L},1)} \right\rangle_{L,\bar{R}} \int d_A^n x\, f_{(\bar{L},2)} = \left\langle \partial^i, 1 \right\rangle_{L,\bar{R}} \int d_A^n x\, f,$$
$$\Rightarrow \quad f_{(\bar{L},1)} \otimes \int d_A^n x\, f_{(\bar{L},2)} = 1 \otimes \int d_A^n x\, f, \tag{313}$$

$$\left\langle \hat{\partial}^i, f_{(L,1)} \right\rangle_{\bar{L},R} \int d_A^n x\, f_{(L,2)} = \left\langle \hat{\partial}^i, 1 \right\rangle_{\bar{L},R} \int d_A^n x\, f,$$
$$\Rightarrow \quad f_{(L,1)} \otimes \int d_A^n x\, f_{(L,2)} = 1 \otimes \int d_A^n x\, f, \tag{314}$$

and

$$\int d_A^n x\, f_{(\bar{R},1)} \left\langle f_{(\bar{R},2)}, \partial^i \right\rangle_{\bar{L},R} = \int d_A^n x\, f \left\langle 1, \partial^i \right\rangle_{\bar{L},R},$$
$$\Rightarrow \quad \int d_A^n x\, f_{(\bar{R},1)} \otimes f_{(\bar{R},2)} = \int d_A^n x\, f \otimes 1, \tag{315}$$

$$\int d_A^n x\, f_{(R,1)} \left\langle f_{(R,2)}, \hat{\partial}^i \right\rangle_{L,\bar{R}} = \int d_A^n x\, f \left\langle 1, \hat{\partial}^i \right\rangle_{L,\bar{R}},$$
$$\Rightarrow \quad \int d_A^n x\, f_{(R,1)} \otimes f_{(R,2)} = \int d_A^n x\, f \otimes 1. \tag{316}$$

To sum up, we have

$$\int_{-\infty}^{+\infty} d_A x^n \, f(x^i) = \int_{-\infty}^{+\infty} d_A x^n \, f(y^j \oplus_{\bar{L}} x^i) = \int_{-\infty}^{+\infty} d_A x^n \, f(x^i \oplus_{\bar{R}} y^j)$$
$$= \int_{-\infty}^{+\infty} d_A x^n \, f(y^j \oplus_L x^i) = \int_{-\infty}^{+\infty} d_A x^n \, f(x^i \oplus_R y^j), \qquad (317)$$

with $A \in \{L, \bar{L}, R, \bar{R}\}$.

Let us return to the property in (301). It tells us that an integral over the whole space is nothing other than a functional which is invariant under translations. For this reason, the problem of constructing an integral over the whole space is equivalent to that of seeking a functional being invariant under translations. To find such a functional it is helpful to realize that all functions subject to

$$\partial^i \triangleright F(x^j) = F(x^j) \triangleleft \partial^i = 0, \quad \text{for all } i, \qquad (318)$$

have to be periodic in the sense

(i) (quantum plane)

$$F(q^{\pm 2} x^1) = F(q^{\pm 2} x^2) = F, \qquad (319)$$

(ii) (three-dimensional Euclidean space)

$$F(q^{\pm 4} x^+) = F(q^{\pm 2} x^3) = F(q^{\pm 4} x^-) = F, \qquad (320)$$

(iii) (four-dimensional Euclidean space)

$$F(q^{\pm 2} x^1) = F(q^{\pm 2} x^2) = F(q^{\pm 2} x^3) = F(q^{\pm 2} x^4) = F, \qquad (321)$$

(iv) (q-deformed Minkowski space)

$$F(q^{\pm 2} r^2) = F(q^{\pm 2} x^+) = F(q^{\pm 2} x^{3/0}) = F(q^{\pm 2} x^-) = F. \qquad (322)$$

These conditions follow from a direct inspection of the results in Ref. [41]. In other words (319)-(322) characterize functions being constant from the point of view provided by q-deformation. For the sake of completeness we

also note that due to (214) and (215) the requirement in (318) is equivalent to

$$F(y^j \oplus_{L/\bar{L}} x^i) = F(x^i), \quad F(x^i \oplus_{R/\bar{R}} y^j) = F(x^i). \tag{323}$$

Now, we are ready to solve the problem of constructing translationally invariant functionals. Functions satisfying the periodicity conditions in (319)-(322) are obtained most easily by means of the Jackson integrals ($a > 0$, $q > 1$)

$$\begin{aligned} F &= (D^i_{q^{\pm a}})^{-1}\big|_0^\infty f = \mp(1-q^{\pm a}) \sum_{k=-\infty}^{\infty} (q^{ak}x^i) f(q^{ak}x^i), \quad x^i > 0, \\ F &= (D^i_{q^{\pm a}})^{-1}\big|_{-\infty}^0 f = \pm(1-q^{\pm a}) \sum_{k=-\infty}^{\infty} (q^{ak}x^i) f(q^{ak}x^i), \quad x^i < 0, \end{aligned} \tag{324}$$

since it holds

$$\begin{aligned} \big((D^i_{q^a})^{-1}\big|_0^\infty f\big)(q^{\pm a}x^i) &= (D^i_{q^a})^{-1}\big|_0^\infty f, \\ \big((D^i_{q^a})^{-1}\big|_{-\infty}^0 f\big)(q^{\pm a}x^i) &= (D^i_{q^a})^{-1}\big|_{-\infty}^0 f. \end{aligned} \tag{325}$$

A short glance at (290)-(293) tells us that our integrals over the whole space are built up out of Jackson integrals. From this observation we see once again that the integrals in (290)-(293) are indeed invariant under translations.

However, there is a point we have to discuss more carefully. Remarkably, the Jackson integrals over the intervals $[\,0, -\infty)$ and $(-\infty, 0\,]$ still depend on the integration variable. Thus, the value of the Jackson integral in (324) is not completely determined. This ambiguity can be removed by setting the value of the integration variable equal to 1 or -1, i.e.

$$(D^i_{q^{\pm a}})^{-1}\big|_{-\infty}^\infty f \equiv \mp(1-q^{\pm a}) \sum_{k=-\infty}^{\infty} q^{ak}\big(f(q^{ak}) + f(-q^{ak})\big). \tag{326}$$

Nevertheless, it arises the question where this ambiguity comes from and what it physically means. First of all, we can say that due to the periodicity of the Jackson integrals in (324) their values can vary within a certain range. So to speak, an infinite Jackson integral cannot be characterized by a certain value, only, but an interval of values. We can believe that this feature expresses the impossibility of making sharp measurements.

There is another very important point which should be clarified. It concerns the behavior of integrals under the action of scaling operators. Up to now we know that the values of infinite Jackson integrals are not uniquely determined, since they depend on a parameter. Important for us is the fact that scaling operators of the form $q^{\alpha\hat{n}_i}$ have an influence on this parameter, when they act on infinite Jackson integrals. This can be seen rather easily by the following calculation:

$$\begin{aligned} q^{\alpha\hat{n}_i} \triangleright (D^i_{q^{\pm a}})^{-1}\big|_0^\infty f &= \mp(1-q^{\pm a}) \sum_{k=-\infty}^{\infty} q^{ak}(q^\alpha x^i) f(q^{ak}(q^\alpha x^i)) \\ &= \big((D^i_{q^{\pm a}})^{-1}\big|_0^\infty f\big)(q^\alpha x^i). \end{aligned} \tag{327}$$

However, if we keep in mind that the parameters of infinite Jackson integrals can take on arbitrary values, we are always allowed to drop scaling operators acting on infinite Jackson integrals.

Now, we can go one step further. The considerations so far show us that right and left invariant functionals are obtained by products of Jackson integrals of the form (326). The integrals in (290)-(293) give examples for such functionals. Due to the identities

$$\begin{aligned} (D^i_{q^a})^{-1}\big|_{-\infty}^\infty f &= q^a (D^i_{q^{-a}})^{-1}\big|_{-\infty}^\infty f, \\ (D^i_{q^a})^{-1}\big|_{-\infty}^\infty q^{c\hat{n}_i} f(x^i) &= q^{-c} q^{c\hat{n}_i} (D^i_{q^a})^{-1}\big|_{-\infty}^\infty f, \end{aligned} \tag{328}$$

which are a direct consequence of the definitions in (324), these integrals are unique up to a global factor if we neglect the parametrization of infinite Jackson integrals and if we do not take into account the appearance of scaling operators.

Up to now we know that integrals over the whole space are invariant under translations and behave like scalars under the action of the algebra describing the quantum symmetry. This way, integrals over the whole space should have trivial braiding. However, a short glance at the L-matrices [cf. (18) and (19)] shows us that the braiding is not only determined by the action of symmetry generators but also by that of certain scaling operators. From the discussion above we know that a scaling operator Λ with $\Lambda \triangleright x^i = q^\alpha x^i$ does not really change a Jackson integral over the intervals $[\,0, -\infty)$ and $(-\infty, 0\,]$. This, in turn, implies that integrals over the whole space remain unchanged under the action of scaling operators. Hence, integrals over the

whole space can indeed be treated as objects having trivial braiding.

Lastly, let us remind of the fact that the explicit form of our integrals depends on the choice for the normal ordering of quantum space coordinates. On this ground, it would be interesting to understand how a change of the underlying normal ordering affects the evaluation of integrals over the whole space. As we know from the discussion in Section 2, there are operators transforming functions to those representing the same quantum space element but for reversed normal ordering. With these operators at hand we show in Appendix B how the expressions for integrals over q-deformed Minkowski space change under reversing the normal ordering. The same method applies to the other quantum spaces we are considering in this article. In this manner we get

(i) (quantum plane)

$$\int_{-\infty}^{+\infty} d^2_{L/\bar{R}}x\, f(x^1,x^2) = \int_{-\infty}^{+\infty} d^2_{L/\bar{R}}x\, q^{-\hat{n}_{x^2}\hat{n}_{x^1}}\tilde{f}(x^2,x^1), \tag{329}$$

$$\int_{-\infty}^{+\infty} d^2_{\bar{L}/R}x\, \tilde{f}(x^2,x^1) = \int_{-\infty}^{+\infty} d^2_{\bar{L}/R}x\, q^{\hat{n}_{x^2}\hat{n}_{x^1}} f(x^1,x^2), \tag{330}$$

(ii) (three-dimensional Euclidean space)

$$\int_{-\infty}^{+\infty} d^3_{L/\bar{R}}x\, f(x^+,x^3,x^-) = \int_{-\infty}^{+\infty} d^3_{L/\bar{R}}x\, q^{2\hat{n}_{x^3}(\hat{n}_{x^+}+\hat{n}_{x^-})}\tilde{f}(x^-,x^3,x^+), \tag{331}$$

$$\int_{-\infty}^{+\infty} d^3_{\bar{L}/R}x\, \tilde{f}(x^-,x^3,x^+) = \int_{-\infty}^{+\infty} d^3_{\bar{L}/R}x\, q^{-2\hat{n}_{x^3}(\hat{n}_{x^+}+\hat{n}_{x^-})} f(x^+,x^3,x^-), \tag{332}$$

(iii) (four-dimensional Euclidean space)

$$\int_{-\infty}^{+\infty} d^4_{\bar{L}/R}x\, f(x^1,x^2,x^3,x^4) =$$

$$\int_{-\infty}^{+\infty} d^4_{\bar{L}/R}x\, q^{-(\hat{n}_{x^2}+\hat{n}_{x^3})(\hat{n}_{x^1}+\hat{n}_{x^4})}\tilde{f}(x^4,x^2,x^3,x^1), \tag{333}$$

$$\int_{-\infty}^{+\infty} d^4_{L/\bar{R}}x\, \tilde{f}(x^4,x^3,x^2,x^1) =$$

$$\int_{-\infty}^{+\infty} d^4_{L/\bar{R}}x\, q^{(\hat{n}_{x^2}+\hat{n}_{x^3})(\hat{n}_{x^1}+\hat{n}_{x^4})}f(x^1,x^2,x^3,x^4), \tag{334}$$

(iv) (q-deformed Minkowski space)

$$\int_{-\infty}^{+\infty} d^4_{\bar{L}/R}x\, f(r^2,x^+,x^{3/0},x^-) =$$

$$\int_{-\infty}^{+\infty} d^4_{\bar{L}/R}x\, q^{2\hat{n}_{x^+}\hat{n}_{x^-}+2\hat{n}_{x^3}(\hat{n}_{x^+}+\hat{n}_{x^-})}\tilde{f}(x^-,x^{3/0},x^+,r^2), \tag{335}$$

$$\int_{-\infty}^{+\infty} d^4_{\bar{L}/R}x\, \tilde{f}(x^-,x^{3/0},x^+,r^2) =$$

$$\int_{-\infty}^{+\infty} d^4_{\bar{L}/R}x\, q^{-2\hat{n}_{x^+}\hat{n}_{x^-}-2\hat{n}_{x^3}(\hat{n}_{x^+}+\hat{n}_{x^-})}f(r^2,x^+,x^{3/0},x^-), \tag{336}$$

where f and $\tilde{f}$ represent the same quantum space element but for different normal orderings.

5.4 Conjugation properties of q-deformed integrals

Now, we turn to the conjugation properties of our integrals. To begin with, let us mention that from an algebraic point of view a consistent conjugation is introduced by

$$\overline{(\partial^i)^{-1}} = (\overline{\partial^i})^{-1}. \tag{337}$$

With this convention the properties in (130) imply for representations of inverse partial derivatives the following identities

$$\begin{aligned}
&\overline{\big((\partial^i)^{-1} \triangleright f(x^j)\big)} = \overline{f(x^j)} \,\bar{\triangleleft}\, (\overline{\partial^i})^{-1}, \quad \overline{\big(f(x^j) \,\bar{\triangleleft}\, (\partial^i)^{-1}\big)} = (\overline{\partial^i})^{-1} \triangleright \overline{f(x^j)}, \\
&\overline{\big((\hat{\partial}^i)^{-1} \,\bar{\triangleright}\, f(x^j)\big)} = \overline{f(x^j)} \triangleleft (\overline{\hat{\partial}^i})^{-1}, \quad \overline{\big(f(x^j) \triangleleft (\hat{\partial}^i)^{-1}\big)} = (\overline{\hat{\partial}^i})^{-1} \,\bar{\triangleright}\, \overline{f(x^j)}.
\end{aligned} \tag{338}$$

Moreover, from (337) it follows that inverse partial derivatives are imaginary, i.e.

$$\overline{(\partial^i)^{-1}} = -(\overline{\partial}_i)^{-1} = -(\partial_i)^{-1} = -g_{ij}(\partial^j)^{-1}. \tag{339}$$

However, for this to be true in general we have to deal with $-\mathrm{i}(\partial^0)^{-1}$ instead of $(\partial^0)^{-1}$ in the case of q-deformed Minkowski space.

In addition to this, we can ask about the conjugation properties of integrals over the whole space. Applying the above results to the expressions in (288) yields

$$\begin{aligned}
\overline{\int_{-\infty}^{+\infty} d_L^n x\, f} &= \lim_{x^{\bar{1}},\ldots,x^{\bar{n}}\to\infty} \overline{(\partial^n)^{-1}\big|_{-\infty}^{x^{\bar{n}}} \ldots (\partial^1)^{-1}\big|_{-\infty}^{x^{\bar{1}}} \triangleright f} \\
&= (-1)^n \lim_{x^1,\ldots,x^n\to\infty} \bar{f} \,\bar{\triangleleft}\, (\partial^{\bar{1}})^{-1}\big|_{-\infty}^{x^1} \ldots (\partial^{\bar{n}})^{-1}\big|_{-\infty}^{x^n} \\
&= (-1)^n \int_{-\infty}^{+\infty} d_{\bar{R}}^n x\, \bar{f},
\end{aligned} \tag{340}$$

$$\begin{aligned}
\overline{\int_{-\infty}^{+\infty} d_R^n x\, f} &= \lim_{x^1,\ldots,x^n\to\infty} \overline{f \triangleleft (\hat{\partial}^{\bar{1}})^{-1}\big|_{-\infty}^{x^1} \ldots (\hat{\partial}^{\bar{n}})^{-1}\big|_{-\infty}^{x^n}} \\
&= (-1)^n \lim_{x^{\bar{1}},\ldots,x^{\bar{n}}\to\infty} (\hat{\partial}^n)^{-1}\big|_{-\infty}^{x^{\bar{n}}} \ldots (\hat{\partial}^1)^{-1}\big|_{-\infty}^{x^{\bar{1}}} \,\bar{\triangleright}\, \bar{f} \\
&= (-1)^n \int_{-\infty}^{+\infty} d_{\bar{L}}^n x\, \bar{f},
\end{aligned} \tag{341}$$

where we used the fact that the metrical factors stemming from (339) cancel

against each other. In very much the same way, we obtain

$$\overline{\int_{-\infty}^{+\infty} d^n_{\bar{L}}x\, f} = (-1)^n \int_{-\infty}^{+\infty} d^n_R x\, \bar{f},$$
$$\overline{\int_{-\infty}^{+\infty} d^n_{\bar{R}}x\, f} = (-1)^n \int_{-\infty}^{+\infty} d^n_L x\, \bar{f}. \tag{342}$$

It is easily seen that under conjugation left integrals are transformed into right ones and vice versa.

If we want to have an integral with

$$\overline{\int f} = \int \bar{f}, \tag{343}$$

it should be clear that we can take the combinations

$$\mathrm{i}^n \int_{-\infty}^{+\infty} d^n_{\bar{L}}x\, f + \mathrm{i}^n \int_{-\infty}^{+\infty} d^n_R x\, f,$$
$$\mathrm{i}^n \int_{-\infty}^{+\infty} d^n_L x\, f + \mathrm{i}^n \int_{-\infty}^{+\infty} d^n_{\bar{R}}x\, f. \tag{344}$$

Notice that for q-deformed Minkowski space integration over the time coordinate requires an additional imaginary factor in the above expressions.

6 Conclusion

Let us end with some comments on what we have done so far. The aim of our program was to provide us with a q-deformed version of analysis on quantum spaces of physical interest, i.e. Manin plane, q-deformed Euclidean space in three or four dimensions, and q-deformed Minkowski space. In our previous work attention was focused on explicit formulae realizing the elements of q-analysis on commutative coordinate algebras. This way, we got multidimensional versions of the well-known q-calculus with Jackson integrals and

Jackson derivatives [62]. In the present article we continued these considerations, but stress was taken on a presentation which reveals the properties of the elements of q-deformed analysis. Remarkably, our reasonings are in complete analogy to the classical situation, which we regain in the undeformed limit as $q \to 1$.

It should be mentioned that in some sense our work is inspired by the so-called star-product formalism. Nevertheless we do not explicitly consider expansions in terms of a deformation parameter, since we try to work with closed expressions, only. In the former literature the problem of formulating physical theories is often attacked by the construction of Hilbert space representations [27, 63–65]. Although this approach is rather rigorous from a mathematical point of view it is not so easy to understand the physical meaning of its results. As was pointed out above, in our framework, however, the relationship to the classical limit is very clear. This observation nourishes the hope that our concept will enable us to develop a q-deformed field theory along the same line of reasonings as its undeformed counterpart.

Acknowledgement

First of all I am very grateful to Eberhard Zeidler for very interesting and useful discussions, special interest in my work, and financial support. Also I wish to express my gratitude to Julius Wess for his efforts, suggestions, and discussions. Furthermore I would like to thank Alexander Schmidt, Fabian Bachmaier, and Florian Koch for useful discussions and their steady support. Finally, I thank Dieter Lüst for kind hospitality.

A Quantum spaces

The coordinates of the two-dimensional q-deformed quantum plane fulfill the relation [51, 70]

$$X^1 X^2 = q X^2 X^1, \tag{345}$$

whereas the quantum metric is given by a matrix ε^{ij} with non-vanishing elements

$$\varepsilon^{12} = q^{-1/2}, \quad \varepsilon^{21} = -q^{1/2}. \tag{346}$$

The inverse of ε^{ij} is given by

$$(\varepsilon^{-1})^{ij} = \varepsilon_{ij} = -\varepsilon^{ij}. \tag{347}$$

In the case of the q-deformed Euclidean space in three dimensions the commutation relations read [18]

$$\begin{aligned} X^3X^+ &= q^2X^+X^3, \\ X^-X^3 &= q^2X^3X^-, \\ X^-X^+ &= X^+X^- + \lambda X^3X^3. \end{aligned} \tag{348}$$

The non-vanishing elements of the quantum metric are

$$g^{+-} = -q, \quad g^{33} = 1, \quad g^{-+} = -q^{-1}. \tag{349}$$

Its inverse is determined by

$$(g^{-1})^{AB} = g_{AB} = g^{AB}. \tag{350}$$

For the four-dimensional q-deformed Euclidean space we have the relations [14, 71]

$$\begin{aligned} X^1X^2 &= qX^2X^1, \\ X^1X^3 &= qX^3X^1, \\ X^3X^4 &= qX^4X^3, \\ X^2X^4 &= qX^4X^2, \\ X^2X^3 &= X^3X^2, \\ X^4X^1 &= X^1X^4 + \lambda X^2X^3. \end{aligned} \tag{351}$$

The metric has as non-vanishing components

$$g^{14} = q^{-1}, \quad g^{23} = g^{32} = 1, \quad g^{41} = q, \tag{352}$$

and it holds

$$(g^{-1})^{ij} = g_{ij} = g^{ij}. \tag{353}$$

In the case of the q-deformed Minkowski space the coordinates obey the relations [14]

$$\begin{aligned} X^\mu X^0 &= X^0X^\mu, \quad \mu \in \{0, +, -, 3\}, \\ X^-X^3 - q^2X^3X^- &= -q\lambda X^0X^-, \\ X^3X^+ - q^2X^+X^3 &= -q\lambda X^0X^+, \end{aligned}$$

$$X^- X^+ - X^+ X^- = \lambda (X^3 X^3 - X^0 X^3), \tag{354}$$

and the non-vanishing components of the metric read

$$\eta^{00} = -1, \quad \eta^{33} = 1, \quad \eta^{+-} = -q, \quad \eta^{-+} = -q^{-1}. \tag{355}$$

Again, we have

$$(\eta^{-1})^{\mu\nu} = \eta_{\mu\nu} = \eta_{\mu\nu}. \tag{356}$$

For other deformations of Minkowski spacetime we refer to Refs. [72–78].

B Integrals for q-deformed Minkowski space

In this appendix we would like to deal with integrals for q-deformed Minkowski space, which from a physical point of view is the most interesting case among q-deformed quantum spaces. Especially, we derive expressions for one-dimensional integrals as well as for integrals over the whole space. In the latter case, we additionally explain how these expressions depend on the choice for the normal ordering.

To begin with, we recall the explicit form of conjugate left representations of partial derivatives [41]:

$$\begin{aligned}
\hat{\partial}^3 \triangleright f(r^2, x^+, x^{3/0}, x^-) &= \underline{D^{3/0}_{q^2} f(q^2 r^2)} \\
&+ q^{-1} \lambda_+ x^{3/0} D^{r^2}_{q^2} f(q^2 x^+) \\
&+ \big(- q^{-2} r^2 (x^{3/0})^{-1} - q^{-4} x^{3/0} \big) D^{r^2}_{q^2} f \\
&+ q^{-2} \lambda_+ x^+ x^- (x^{3/0})^{-1} D^{r^2}_{q^2} f(q^2 x^{3/0}) \\
&- q \lambda_+^{-1} \lambda x^{3/0} D^+_{q^2} D^-_{q^2} f(q^2 r^2),
\end{aligned} \tag{357}$$

$$\begin{aligned}
\hat{\partial}^+ \triangleright f(r^2, x^+, x^{3/0}, x^-) &= \underline{-q D^-_{q^2} f(q^2 r^2)} \\
&+ q^{-1} \lambda_+ x^+ D^{r^2}_{q^2} f,
\end{aligned} \tag{358}$$

$$\begin{aligned}
\hat{\partial}^- \triangleright f(r^2, x^+, x^{3/0}, x^-) &= \underline{-q^{-1} D^+_{q^2} f} \\
&+ q^{-1} \lambda_+ x^- D^{r^2}_{q^2} f(q^2 x^+, q^2 x^{3/0}) \\
&+ q^{-2} \lambda (x^{3/0})^2 D^+_{q^2} D^{r^2}_{q^2} f(q^2 x^+, q^2 x^-)
\end{aligned}$$

$$- \; q^{-1}\lambda^2 (x^{3/0})^2 x^- D^+_{q^2} D^-_{q^2} D^{r^2}_{q^2} f(q^2 x^+), \tag{359}$$

$$\begin{aligned}
\hat{\partial}^2 \bar{\triangleright} f(r^2, x^+, x^{3/0}, x^-) &= \underline{q\lambda_+^3 D^{r^2}_{q^2} f(q^2 x^+, q^2 x^{3/0}, q^2 x^-)} \\
&+ q^3 \lambda_+^3 r^2 (D^{r^2}_{q^2})^2 f(q^2 x^+, q^2 x^{3/0}, q^2 x^-) \\
&+ \; (q^{-1}\lambda_+)^2 x^- D^-_{q^2} D^{r^2}_{q^2} f(q^2 x^+, q^2 x^{3/0}) \\
&+ \; (q^{-1}\lambda_+)^2 x^{3/0} D^{3/0}_{q^2} D^{r^2}_{q^2} f(q^2 x^+) \\
&- \; \lambda_+ D^+_{q^2} D^-_{q^2} f(q^2 r^2) + (q^{-1}\lambda_+)^2 x^+ D^+_{q^2} D^{r^2}_{q^2} f,
\end{aligned} \tag{360}$$

where $\lambda_+ \equiv q + q^{-1}$.

Next, we wish to apply the result of (244) to the above representations. To this end, we first have to identify the expressions for the $\hat{\partial}^i_{\rm cl}$. They have to be invertible operators that do not vanish in the classical limit. In this manner, we choose for the $\hat{\partial}^i_{\rm cl}$ the expressions we have underlined in (357)-(360). We find as corresponding inverse operators

$$\begin{aligned}
(\hat{\partial}^3_{\rm cl})^{-1} \bar{\triangleright} f &= (D^{3/0}_{q^2})^{-1} f(q^{-2} r^2), \\
(\hat{\partial}^+_{\rm cl})^{-1} \bar{\triangleright} f &= -q^{-1} (D^-_{q^2})^{-1} f(q^{-2} r^2), \\
(\hat{\partial}^-_{\rm cl})^{-1} \bar{\triangleright} f &= -q (D^+_{q^2})^{-1} f, \\
(\hat{\partial}^{r^2}_{\rm cl})^{-1} \bar{\triangleright} f &= q^{-1} \lambda_+^{-3} (D^{r^2}_{q^2})^{-1} f(q^{-2} x^+, q^{-2} x^{3/0}, q^{-2} x^-).
\end{aligned} \tag{361}$$

Consequently, the terms being not underlined contribute to $\hat{\partial}^i_{\rm cor}$. Inserting the explicit form of $(\hat{\partial}^i_{\rm cl})^{-1}$ and $\hat{\partial}^i_{\rm cor}$ into (244) then yields the wanted expressions for the integral operator $(\hat{\partial}^i)^{-1}$. We leave the details to the reader and write down the result for $(\hat{\partial}^+)^{-1}$ as an example:

$$\begin{aligned}
(\hat{\partial}^+)^{-1} \bar{\triangleright} f = -q^{-1} \sum_{k=0}^{\infty} &(q^{-2}\lambda_+)^k q^{k(k+1)} (x^+)^k \\
&\times (D^{r^2}_{q^2})^k (D^-_{q^2})^{-(k+1)} f(q^{-2(k+1)} r^2).
\end{aligned} \tag{362}$$

Now, we come to integrals over the whole Minkowski-space. They can be introduced by

$$\int d^4_{\bar{L}} x \, f(x^i) \equiv (\hat{\partial}^2)^{-1}\big|^{\infty}_{-\infty} (\hat{\partial}^+)^{-1}\big|^{\infty}_{-\infty} (\hat{\partial}^3)^{-1}\big|^{\infty}_{-\infty} (\hat{\partial}^-)^{-1}\big|^{\infty}_{-\infty} \bar{\triangleright} f. \tag{363}$$

It is our aim to rewrite the last expression in terms of Jackson integrals. To this end we first try to simplify the formulae for the one-dimensional integrals. In the case of $(\hat{\partial}^-)^{-1}$, for example, we can proceed as follows:

$$
\begin{aligned}
(\hat{\partial}^-)^{-1}\big|_{-\infty}^{\infty} \bar{\triangleright}\, f &= \sum_{k=0}^{\infty} (-1)^k \left[(\hat{\partial}^-_{\mathrm{cl}})^{-1}\big|_{-\infty}^{\infty} \hat{\partial}^-_{\mathrm{cor}}\right]^k (\hat{\partial}^-_{\mathrm{cl}})^{-1}\big|_{-\infty}^{\infty} f \\
&= -q \sum_{k=0}^{\infty} \Big[q(D^+_{q^2})^{-1}\big|_{-\infty}^{\infty} \Big(q^{-1}\lambda_+ x^- D^{r^2}_{q^2} q^{2(\hat{n}_{x^+}+\hat{n}_{x^{3/0}})} \\
&\qquad + q^{-2}\lambda (x^{3/0})^2 D^+_{q^2} D^{r^2}_{q^2} q^{2(\hat{n}_{x^+}+\hat{n}_{x^-})} \\
&\qquad - q^{-1}\lambda^2 (x^{3/0})^2 x^- D^+_{q^2} D^-_{q^2} D^{r^2}_{q^2} q^{2\hat{n}_{x^+}} \Big) \Big]^k (D^+_{q^2})^{-1}\big|_{-\infty}^{\infty} f \\
&= -q \sum_{k=0}^{\infty} \left[\lambda_+ x^- (D^+_{q^2})^{-1}\big|_{-\infty}^{\infty} D^{r^2}_{q^2} q^{2(\hat{n}_{x^+}+\hat{n}_{x^{3/0}})} \right]^k (D^+_{q^2})^{-1}\big|_{-\infty}^{\infty} f \\
&= -q \sum_{k=0}^{\infty} q^{k(k+1)} (\lambda_+ x^- D^{r^2}_{q^2})^k (D^+_{q^2})^{-(k+1)}\big|_{-\infty}^{\infty} f(q^{2k}x^+, q^{2k}x^{3/0}). \qquad (364)
\end{aligned}
$$

The first equality is the formula from (244), for the second equality we insert the explicit form of $(\hat{\partial}^-_{\mathrm{cl}})^{-1}$ and $\hat{\partial}^-_{\mathrm{cor}}$, while the third equality uses

$$
\begin{aligned}
D^i_{q^a}(D^i_{q^a})^{-1}\big|_{-\infty}^{\infty} f &= \frac{(D^i_{q^a})^{-1}\big|_{-\infty}^{\infty} f - \big((D^i_{q^a})^{-1}\big|_{-\infty}^{\infty} f\big)(q^a x^i)}{(1-q^a)x^i} \\
&= \frac{(D^i_{q^a})^{-1}\big|_{-\infty}^{\infty} f - (D^i_{q^a})^{-1}\big|_{-\infty}^{\infty} f}{(1-q^a)x^i} \\
&= 0. \qquad (365)
\end{aligned}
$$

In very much the same way we can show that

$$
\begin{aligned}
&(\hat{\partial}^2)^{-1}\big|_{-\infty}^{\infty} \bar{\triangleright}\, f = \\
&\sum_{k=0}^{\infty} q^{(\frac{3}{2}k-1)(k+1)} \lambda_+^{-2k-3} (D^+_{q^2} D^-_{q^2})^k \\
&\times (D^{r^2}_{q^2})^{-(k+1)}\big|_{-\infty}^{\infty} f(q^{2k}r^2, q^{-2(k+1)}x^+, q^{-2(k+1)}x^{3/0}, q^{-2(k+1)}x^-). \qquad (366)
\end{aligned}
$$

Putting everything together we obtain

$$
\begin{aligned}
&(\hat{\partial}^{+})^{-1}\big|_{-\infty}^{\infty}(\hat{\partial}^{3})^{-1}\big|_{-\infty}^{\infty}(\hat{\partial}^{-})^{-1}\big|_{-\infty}^{\infty}(\hat{\partial}^{2})^{-1}\big|_{-\infty}^{\infty}\triangleright f = \\
&\quad\sum_{j=0}^{\infty}(q^{-2}\lambda_{+})^{j}q^{j(j+1)}(x^{+})^{j}(D_{q^2}^{r^2})^{j}(D_{q^2}^{-})^{-(j+1)}\big|_{-\infty}^{\infty}q^{-2(j+1)\hat{n}_{r^2}} \\
&\quad\times\sum_{k=0}^{\infty}(-1)^{k}\Big[(D_{q^2}^{3/0})^{-1}\big|_{-\infty}^{\infty}q^{-2\hat{n}_{r^2}}\Big(q^{-1}\lambda_{+}x^{3/0}D_{q^2}^{r^2}q^{2\hat{n}_{x^{+}}} \\
&\qquad+\big(-q^{-2}r^{2}(x^{3/0})^{-1}-q^{-4}x^{3/0}\big)D_{q^2}^{r^2} \\
&\qquad+q^{-2}\lambda_{+}x^{+}x^{-}(x^{3/0})^{-1}D_{q^2}^{r^2}q^{-2\hat{n}_{x^{3/0}}} \\
&\qquad-q\lambda_{+}^{-1}\lambda x^{3/0}D_{q^2}^{+}D_{q^2}^{-}q^{2\hat{n}_{r^2}}\Big)\Big]^{k}(D_{q^2}^{3/0})^{-1}\big|_{-\infty}^{\infty}q^{-2\hat{n}_{r^2}} \\
&\quad\times\sum_{l=0}^{\infty}q^{l(l+1)}(\lambda_{+}x^{-}D_{q^2}^{r^2})^{l}(D_{q^2}^{+})^{-(l+1)}\big|_{-\infty}^{\infty}q^{2l(\hat{n}_{x^{+}}+\hat{n}_{x^{3/0}})}f \\
&\quad\times\sum_{m=0}^{\infty}q^{(\frac{3}{2}m-1)(m+1)}\lambda_{+}^{-2m-3}(D_{q^2}^{+}D_{q^2}^{-})^{m} \\
&\quad\times(D_{q^2}^{r^2})^{-(m+1)}\big|_{-\infty}^{\infty}f(q^{2m}r^{2},q^{-2(m+1)}x^{+},q^{-2(m+1)}x^{3/0},q^{-2(m+1)}x^{-}) \\
&=-q^{-1}\lambda_{+}^{-3}(D_{q^2}^{-})^{-1}\big|_{-\infty}^{\infty}q^{-2\hat{n}_{r^2}}(D_{q^2}^{3/0})^{-1}\big|_{-\infty}^{\infty}q^{-2\hat{n}_{r^2}} \\
&\quad\times(D_{q^2}^{+})^{-1}\big|_{-\infty}^{\infty}(D_{q^2}^{r^2})^{-1}\big|_{-\infty}^{\infty}q^{-2(\hat{n}_{x^{+}}+\hat{n}_{x^{3/0}}+\hat{n}_{x^{-}})}f.
\end{aligned}
\tag{367, 368}
$$

For the first identity the expressions we have derived so far are substituted for the one-dimensional integrals. The second equality makes use of (365) and

$$
(D_{q^a}^{i})^{-1}\big|_{-\infty}^{\infty}(D_{q^a}^{i})^{n}f=\lim_{x^{i}\to\infty}(D_{q^a}^{i})^{n-1}f-\lim_{x^{i}\to-\infty}(D_{q^a}^{i})^{n-1}f=0. \tag{369}
$$

Notice that the last result only holds if we assume $f\in\widetilde{\mathcal{A}}$ [cf. (257)].

Lastly, we would like to discuss how the expression in (368) changes under reversing the normal ordering. To this end we need an operator that transforms a function representing an element of q-deformed Minkowski space for ordering $X^{-}X^{3/0}X^{+}\hat{r}^{2}$ to a function representing the same quantum space element but now for reversed ordering $\hat{r}^{2}X^{+}X^{3/0}X^{-}$. Its explicit form reads [38]

$$
f(r^{2},x^{+},x^{3/0},x^{-})=\hat{U}^{-1}\big(f(x^{-},x^{3/0},x^{+},r^{2})\big)
$$

$$= \sum_{i,j=0}^{\infty} (\lambda\lambda_+^{-1})^i \frac{(-1)^j q^{-(i-k)-k^2}}{[[k]]_{q^2}![[i-k]]_{q^2}!}$$
$$\times r^{2(i-k)} \big(x^{3/0}\big)^{2k} q^{2\hat{n}_{x^+}\hat{n}_{x^-} + (\hat{n}_{x^+}+\hat{n}_{x^-})(2\hat{n}_{x^{3/0}}+i)+2\hat{n}_{x^{3/0}}i}$$
$$\times \big((D_{q^2}^+ D_{q^2}^-)^i \tilde{f}\big)(q^{i-2k}x^-, x^{3/0}, q^{i-2k}x^+, r^2). \qquad (370)$$

As a next step we take the integral of $f = \hat{U}^{-1}\tilde{f}$ over the whole Minkowski space. By virtue of (368) and (369) we finally arrive at

$$\int d^4_{\tilde{L}} x\, f = \int d^4_{\tilde{L}} x\, q^{2\hat{n}_{x^+}\hat{n}_{x^-} + 2(\hat{n}_{x^+}+\hat{n}_{x^-})\hat{n}_{x^{3/0}}}\, \tilde{f}, \qquad (371)$$

i.e. the terms with $i > 0$ in (370) do not contribute to the integral.

References

[1] J. Schwinger, *Quantum Electrodynamics*, Dover, New York, 1958.

[2] S. Weinberg, *Gravitation and Cosmology,* Wiley, New York, 1972.

[3] D. C. Cassidy, *Uncertainty: The Life and Science of Werner Heisenberg*, Freeman, New York, 1992.

[4] W. Heisenberg, *Über die in der Theorie der Elementarteilchen auftretende universelle Länge,* Ann. Phys. **32** (1938) 20.

[5] I. M. Gelfand and M. A. Naimark, *On the embedding of normed linear rings into the ring of operators in Hilbert space*, Math. Sbornik **12** (1943) 197.

[6] H. Hopf, *Über die Topologie der Gruppenmannigfaltigkeiten und ihre Verallgemeinerungen*, Ann. Math. **42** (1941) 22.

[7] P. P. Kulish and N. Y. Reshetikin, *Quantum linear problem for the Sine-Gordon equation and higher representations,* J. Sov. Math. **23** (1983) 2345.

[8] S. L. Woronowicz, *Compact matrix pseudogroups,* Commun. Math. Phys. **111** (1987) 613.

[9] V. G. Drinfeld, *Hopf algebras and the quantum Yang-Baxter equation,* Sov. Math. Dokl. **32** (1985) 254.

[10] M. Jimbo, *A q-analogue of U(g) and the Yang-Baxter equation,* Lett. Math. Phys. **10** (1985) 63.

[11] V. G. Drinfeld, *Quantum groups*, in A. M. Gleason, ed., Proceedings of the International Congress of Mathematicians, Amer. Math. Soc., 798 (1986).

[12] N. Yu. Reshetikhin, L. A. Takhtadzhyan and L. D. Faddeev, *Quantization of Lie Groups and Lie Algebras*, Leningrad Math. J. **1** (1990) 193.

[13] M. Takeuchi, *Matrix Bialgebras and Quantum Groups,* Israel J. Math. **72** (1990) 232.

[14] U. Carow-Watamura, M. Schlieker, M. Scholl and S. Watamura, *Tensor Representations of the Quantum Group $SL_q(2)$ and Quantum Minkowski Space,* Z. Phys. C **48** (1990) 159.

[15] P. Podleś and S. L. Woronowicz, *Quantum Deformation of Lorentz Group*, Commun. Math. Phys. **130** (1990) 381.

[16] W. B. Schmidke, J. Wess and B. Zumino, *A q-deformed Lorentz Algebra in Minkowski phase space,* Z. Phys. C **52** (1991) 471.

[17] S. Majid, *Examples of braided groups and braided matrices,* J. Math. Phys. **32** (1991) 3246.

[18] A. Lorek, W. Weich and J. Wess, *Non-commutative Euclidean and Minkowski Structures,* Z. Phys. C **76** (1997) 375, [q-alg/9702025].

[19] J. Wess and B. Zumino, *Covariant differential calculus on the quantum hyperplane*, Nucl. Phys. B. Suppl. **18** (1991) 302.

[20] U. Carow-Watamura, M. Schlieker and S. Watamura, *$SO_q(N)$-covariant differential calculus on quantum space and deformation of Schrödinger equation,* Z. Phys. C **49** (1991) 439.

[21] X. C. Song, *Covariant differential calculus on quantum minkowski space and q-analog of Dirac equation,* Z. Phys. C **55** (1992) 417.

[22] O. Ogievetsky, W. B. Schmidke, J. Wess and B. Zumino, *q-Deformed Poincaré Algebra,* Commun. Math. Phys. **150** (1992) 495.

[23] S. Majid, *Braided momentum in the q-Poincaré group*, J. Math. Phys. **34** (1993) 2045.

[24] M. Fichtenmüller, A. Lorek and J. Wess, *q-deformed Phase Space and its Lattice Structure,* Z. Phys. C **71** (1996) 533, [hep-th/9511106].

[25] J. Wess, *q-Deformed phase space and its lattice structure,* Int. J. Mod. Phys. A **12** (1997) 4997.

[26] B. L. Cerchiai, R. Hinterding, J. Madore and J. Wess, *A Calculus Based on a q-deformed Heisenberg Algebra,* Eur. Phys. J. C **8** (1999) 547, [math-ph/9809160].

[27] B. L. Cerchiai and J. Wess, *q-Deformed Minkowski Space based on a q-Lorentz Algebra,* Eur. Phys. J. C **5** (1998) 553, [math.qa/9801104].

[28] H. Grosse, C. Klimčik and P. Prešnajder, *Towards finite quantum field theory in non-commutative geometry,* Int. J. Theor. Phys. **35** (1996) 231, [hep-th/9505175].

[29] S. Majid, *On the q-regularisation,* Int. J. Mod. Phys. A **5** (1990) 4689.

[30] R. Oeckl, *Braided Quantum Field Theory,* Commun. Math. Phys. **217** (2001) 451.

[31] C. Blohmann, *Free q-deformed relativistic wave equations by representation theory*, Eur. Phys. J. C **30** (2003) 435, [hep-th/0111172].

[32] J. Wess, *q-deformed Heisenberg Algebras*, in H. Gausterer, H. Grosse and L. Pittner, eds., Proceedings of the 38. Internationale Universitätswochen für Kern- und Teilchenphysik, no. 543 in Lect. Notes in Phys., Springer-Verlag, Schladming (2000), [math-ph/9910013].

[33] S. Majid, *Representations, duals and quantum doubles of monoidal categories,* Suppl. Rend. Circ. Mat. Palermo, Ser. II, **26** (1991) 197.

[34] S. Majid, *Algebras and Hopf Algebras in Braided Categories,* Lec. Notes Pure Appl. Math. **158** (1994) 55.

[35] S. Majid, *Beyond Supersymmetry and Quantum Symmetry (an itroduction to braided groups and braided matrices)*, in M. L. Ge and H. J. de Vega, eds., Quantum Groups, Integrable Statistical Models and Knot Theory, World Scientific, 231 (1993).

[36] S. Majid, *Introduction to Braided Geometry and q-Minkowski Space,* preprint (1994), [hep-th/9410241].

[37] H. Wachter and M. Wohlgenannt, $*$*-Products on quantum spaces,* Eur. Phys. J. C **23** (2002) 761, [hep-th/0103120].

[38] C. Bauer and H. Wachter, *Operator representations on quantum spaces,* Eur. Phys. J. C **31** (2003) 261, [math-ph/0201023].

[39] H. Wachter, *q-Integration on quantum spaces,* Eur. Phys. J. C **32** (2004) 281, [hep-th/0206083].

[40] H. Wachter, *q-Exponentials on quantum spaces,* Eur. Phys. J. C **37** (2004) 379, [hep-th/0401113].

[41] H. Wachter, *q-Translations on quantum spaces,* preprint, [hep-th/0410205].

[42] H. Wachter, *Braided products for quantum spaces,* preprint, [math-ph/0509018].

[43] D. Mikulovic, A. Schmidt and H. Wachter, *Grassmann variables on quantum spaces,* Eur. Phys. J. C **45** (2006) 529, [hep-th/0407273].

[44] A. Schmidt, H. Wachter, *Superanalysis on quantum spaces*, JHEP **0601** (2006) 84, [hep-th/0411180].

[45] S. Majid, *Braided geometry: A new approach to q-deformations,* in R. Coquereaux et al., eds., First Caribbean Spring School of Mathematics and Theoretical Physics, World Scientific, Guadeloupe, 190 (1993).

[46] S. Majid, *Free braided differential calculus, braided binomial theorem and the braided exponential map,* J. Math. Phys. **34** (1993) 4843.

[47] S. Majid, *Foundations of Quantum Group Theory,* University Press, Cambridge (1995).

[48] O. Ogievetsky and B. Zumino, *Reality in the differential calculus on q-Euclidean spaces*, Lett. Math. Phys. **25** (1992) 121.

[49] S. Majid, $*$*-structures on braided spaces*, J. Math. Phys. **36** (1995) 4436.

[50] S. Majid, *Quasi-$*$-structure on q-Poincaré algebras*, J. Geom. Phys. **22** (1997) 14.

[51] Y. I. Manin, *Quantum Groups and Non-Commutative Geometry*, Centre de Recherche Mathématiques, Montreal (1988).

[52] F. Bayen, M. Flato, C. Frønsdal, A. Lichnerowicz and D. Sternheimer, *Deformation theory and quantization. I. Deformations of symplectic structures*, Ann. Phys. **111** (1978) 61.

[53] J. E. Moyal, *Quantum mechanics as a statistical theory,* Proc. Camb. Phil. Soc. **45** (1949) 99.

[54] J. Madore, S. Schraml, P. Schupp and J. Wess, *Gauge Theory on Non-commutative Spaces,* Eur. Phys. J. C **16** (2000) 161, [hep-th/0001203].

[55] F. H. Jackson, *On q-functions and a certain difference operator,* trans. Roy. Edin. **46** (1908) 253.

[56] A. Klimyk and K. Schmüdgen, *Quantum Groups and their Representations,* Springer Verlag, Berlin (1997).

[57] S. Majid, *Quantum and braided linear algebra*, J. Mat. Phys. **34** (1993) 1176.

[58] U. Meyer, *q-Lorentz group and braided coaddition on q-Minkowski space*, Commun. Math. Phys. **168** (1995) 249.

[59] S. M. Lane, *Categories for the Working Mathematician*, Springer Verlag, 1974.

[60] M. Chaichian and A. P. Demichev, *Introduction to Quantum Groups*, World Scientific, Singapore, 1996.

[61] F. H. Jackson, *q-Integration,* Proc. Durham Phil. Soc. **7** (1927) 182.

[62] V. Kac and P. Cheung, *Quantum Calculus,* Springer Verlag, Berlin (2000).

[63] G. Fiore, *The $SO_q(N)$-symmmetric harmonic oszillator on the quantum Euclidean space R_q^N and its Hilbert space structure*, Int. J. Mod. Phys. A **8** (1993) 4679.

[64] W. Zippold, *Hilbertspace Representations of an Algebra of Observables for q-deformed Relativistic Quantum Mechanics*, Z. Phys. C **67** (1995) 681.

[65] F. Bachmaier, *The free particle on q-Minkowski space*, Doktorarbeit, Ludwig-Maximilians-Universität München, Fakultät für Physik (2003).

[66] C. Chryssomalakos, P. Schupp and P. Watts, *Translations, Integrals And Fourier Transforms In The Quantum Plane*, in A. Ali, J. Ellis and S. Randjbar-Daemi, eds., Salamfestschrift, Proceedings of the Conference on Highlights of Particle and Condensed Matter Physics, ICTP, Trieste (1993).

[67] A. Kempf and S. Majid, *Algebraic q-integration and Fourier theory on quantum and braided spaces,* J. Math. Phys. **35** (1994) 6802.

[68] C. Chryssomalakos, *Remarks On Quantum Integration*, preprint, ENSLAPP-A-562/95, [q-alg/9601014].

[69] H. Steinacker, *Integration on quantum Euclidean Space and sphere,* J. Math. Phys. **37** (1996) 4738.

[70] M. Schlieker and W. Scholl, *Spinor calculus for quantum groups*, Z. Phys. C **47** (1990) 625.

[71] H. Ocambo, *$SO_q(4)$ quantum mechanics*, Z. Phys. C **70** (1996) 525.

[72] J. Lukierski, A. Nowicki and H. Ruegg, *New Quantum Poincaré Algebra and κ-deformed Field Theory,* Phys. Lett. B **293** (1992) 344.

[73] L. Castellani, *Differential Calculus on $ISO_q(N)$, Quantum Poincaré Algebra and q-Gravity,* preprint, [hep-th/9312179].

[74] V. K. Dobrev, *New q-Minkowski space-time and q-Maxwell equations hierarchy from q-conformal invariance,* Phys. Lett. B **341** (1994) 133.

[75] M. Chaichian and A. P. Demichev, *Quantum Poincaré group without dilatation and twisted classical algebra*, J. Math. Phys. **36** (1995) 398.

[76] S. Doplicher, K. Fredenhagen and J. E. Roberts, *The Quantum Structure of Space-Time at the Planck Scale and Quantum fields*, Commun. Math. Phys. **172** (1995) 187.

[77] M. Chaichian, P. P. Kulish, K. Nishijima and A. Tureanu, *On a Lorentz-Invariant Interpretation of Noncommutative Space-Time and its Implications on Noncommutative QFT,* Phys. Lett. B **604** (2004) 98, [hep-th/0408062].

[78] F. Koch and E. Tsouchnika, *Construction of θ-Poincaré Algebras and their invariants on* $\mathcal{M}_\theta$, Nucl. Phys. B **717** (2005) 387, [hep-th/0409012].